



=>



chain nodes :

12 13 14 15

ring nodes :

1 2 3 4 5 6 7 8 9

chain bonds :

3-12 8-13 12-14 13-15

ring bonds :

1-2 1-5 1-9 2-3 3-4 4-5 5-6 6-7 7-8 8-9

exact/norm bonds :

1-2 1-5 1-9 2-3 3-4 3-12 4-5 5-6 6-7 7-8 8-9 8-13 12-14 13-15

G1:C,N

Match level :

1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:Atom 8:Atom 9:Atom 12:Atom 13:Atom 14:Atom 15:Atom

L1 STRUCTURE UPLOADED

=> s l1 sss sam
SAMPLE SEARCH INITIATED 10:36:59 FILE 'REGISTRY'
SAMPLE SCREEN SEARCH COMPLETED - 11451 TO ITERATE
17.5% PROCESSED 2000 ITERATIONS
INCOMPLETE SEARCH (SYSTEM LIMIT EXCEEDED)
SEARCH TIME: 00.00.01

1 ANSWERS

FULL FILE PROJECTIONS: ONLINE **COMPLETE**
BATCH **COMPLETE**
PROJECTED ITERATIONS: 222608 TO 235432
PROJECTED ANSWERS: 1 TO 257

L2 1 SEA SSS SAM L1

=> s l1 sss full
FULL SEARCH INITIATED 10:37:08 FILE 'REGISTRY'
FULL SCREEN SEARCH COMPLETED - 228722 TO ITERATE

100.0% PROCESSED 228722 ITERATIONS

265 ANSWERS

SEARCH TIME: 00.00.03

L3 265 SEA SSS FUL L1

-> d scan L3

L3 265 ANSWERS REGISTRY COPYRIGHT 2008 ACS on STN

Author/Inventor

Acetamide, N-[[2-(4-methylphenyl)-6-(1-pyrrolidinyl)imidazo[1,2-b]pyridazin-3-yl]methyl]-

L3 265 ANSWERS REGISTRY COPYRIGHT 2008 ACS on STN

Author/Inventor

Imidazo[1,2-b]pyridazine, 2-[4-{1-methyl-5-(trifluoromethyl)-1H-pyrazol-3-yl}phenyl]-6-(2H-tetrazol-5-yl)-

L3 265 ANSWERS REGISTRY COPYRIGHT 2008 ACS on STN

Author/Inventor

Benzenecarboximidamide, 4,4'-imidazo[1,2-a]pyridine-2,6-diylbis[N-hydroxy- (9CI)

L3 265 ANSWERS REGISTRY COPYRIGHT 2008 ACS on STN

Author/Inventor

Imidazo[1,2-a]pyridine, 6-(2-naphthalenyl)-2-[4-{10-(2-naphthalenyl)-9-anthracenyl}phenyl]-

-> d his

(FILE 'HOME' ENTERED AT 10:36:15 ON 09 JAN 2008)

FILE 'REGISTRY' ENTERED AT 10:36:36 ON 09 JAN 2008

L1 STRUCTURE UPLOADED

L2 1 S L1 SSS SAM

L3 265 S L1 SSS FULL

FILE 'CAPLUS' ENTERED AT 10:37:59 ON 09 JAN 2008

-> s l3

L4 65 L3

-> s l4 and py<=2003

23975208 PY<=2003

L5 42 L4 AND PY<=2003

=> s l5 and electroluminescent

68951 ELECTROLUMINESCENT

6 ELECTROLUMINESCENTS

68953 ELECTROLUMINESCENT

(ELECTROLUMINESCENT OR ELECTROLUMINESCENTS)

L6 1 L5 AND ELECTROLUMINESCENT

-> d l6

L6 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Organic electroluminescent device

Author/Inventor

Nakatsuka, Masakatsu; Shimamura, Takehiko

Patent Assignee/Corporate Source

Mitsui Chemicals Inc., Japan

Source

Jpn. Kokai Tokkyo Koho, 43 pp. CODEN: JKXXAF

Document Type

Patent

Language

Japanese

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2001035664	A	20010209	JP 1999-206147	19990721

Patent Number (1)

JP 2001035664

Kind Code (1)

A

Patent Publication Date (1)
20010209
Application Number (1)
JP 1999-206147
Application Date (1)
19990721
Priority Patent Number (1)
JP 1999-206147
Priority Patent Publication Date (1)
19990721

=> s 15 not 16
L7 41 L5 NOT L6

=> s 17 and electron
1449299 ELECTRON
272382 ELECTRONS
1535444 ELECTRON
(ELECTRON OR ELECTRONS)
L8 1 L7 AND ELECTRON

=> d 18 ibib abs

L8 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2008 ACS on STN
Title

Imidazo[1,2-a]pyridines. II. Ozonolysis of imidazo[1,2-a]pyridines and synthesis of cardiotonic agents

Author/Inventor

Yamanaka, Motosuke; Suda, Shinji; Yoneda, Naoki; Ohara, Hideto

Patent Assignee/Corporate Source

Eisai Tsukuba Res. Lab., Tsukuba, 300-26, Japan

Source

Chemical & Pharmaceutical Bulletin (1992), 40(3), 666-74 CODEN: CPBTAL; ISSN: 0009-2363

Document Type

Journal

Language

English

Abstract

The metabolite of loproline (E-1020) in dogs, 5-(2-aminopyridin-5-yl)-1,2-dihydro-6-methyl-2-oxo-3-pyridinecarboxitrile (I), was prepared via ozonolysis of imidazo[1,2-a]pyridinylpyridines, e.g., II, and evaluated for pos. inotropic activity. Its potency was less than that of loproline and milrinone. Among compds. related to loproline which were synthesized using the versatile intermediates III (R = Me, PhCH₂), obtained during the preparation of I, only 5-(2-aminoimidazo[1,2-a]pyridin-6-yl)-1,2-dihydro-6-methyl-2-oxo-3-pyridinecarboxitrile (IV) retained the activity of loproline. Electron-withdrawing substituents at the 2-position of imidazo[1,2-a]pyridine reduced the activity of the parent compound. The ozonolysis of imidazo[1,2-a]pyridine derivs. under neutral conditions afforded 2-acylamino-pyridine derivs. in a 30-55% yield independent of the substituents at the 2-position of imidazo[1,2-a]pyridine. It is possible to use imidazo[1,2-a]pyridines as protected 2-aminopyridines, and 2,3-unsubstituted imidazo[1,2-a]pyridines are convenient for that purpose from the viewpoint of ease of preparation of the starting material.

=> s 17 not 18
L9 40 L7 NOT L8

=> s 19 and transport
767717 TRANSPORT
6560 TRANSPORTS
770375 TRANSPORT
(TRANSPORT OR TRANSPORTS)
L10 0 L9 AND TRANSPORT

=> s 110 and conduct?
1080537 CONDUCT?
354282 COND
4753 CONDS
355482 COND
(COND OR CONDS)
1209464 CONDUCT?
(CONDUCT? OR COND)
L11 0 L10 AND CONDUCT?

=> s 19 and conduct?
1080537 CONDUCT?
354282 COND
4753 CONDS
355482 COND

{COND OR CONDS}
1209464 CONDUCT?
{CONDUCT? OR COND}
L12 1 L9 AND CONDUCT?

=> d 112 ibib abs

L12 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Electrophotographic photoreceptor using novel azo-type photoconductive substance

Author/Inventor

Ito, Akira; Horiuchi, Tamotsu

Patent Assignee/Corporate Source

Mitsubishi Paper Mills Ltd, Japan

Source

Jpn. Kokai Tokkyo Koho, 18 pp. CODEN: JKXXAF

Document Type

Patent

Language

Japanese

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 08101520	A	19960416	JP 1994-237621	19940930

Patent Number (1)

JP 08101520

Kind Code (1)

A

Patent Publication Date (1)

19960416

Application Number (1)

JP 1994-237621

Application Date (1)

19940930

Priority Patent Number (1)

JP 1994-237621

Priority Patent Publication Date (1)

19940930

Abstract

The title photoreceptor comprises a conductive support laminated with a photosensitive layer containing an azo compound I [R1 = H, halo, CN, (substituted) alkyl, aryl; R2-4 = H, halo, (substituted) alkyl, alkoxy, aryl, heterocyclyl; Ar = (substituted) arylene; n = 0, 1; Cp = coupler residue]. The photoreceptor shows high photosensitivity and durability in repeated used. Thus, an Al vapor-deposited polyester film was coated with a charge-generating layer containing II and a charge-transporting layer containing a hydrazone compound to give a photoreceptor.

=> s 19 not 112
L13 39 L9 NOT L12

=> d 113 1-39 ibib abs

L13 ANSWER 1 OF 39 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Synthesis and antimicrobial evaluation of some new 1,2,4-triazine derivatives

Author/Inventor

El-Ansary, A. K.

Patent Assignee/Corporate Source

Organic Chemistry Department, Faculty of Pharmacy, Cairo University, Cairo, Egypt

Source

Bulletin of the Faculty of Pharmacy (Cairo University) (2002), 40(3), 97-108 CODEN: BFPHA8; ISSN: 1110-0931

Document Type

Journal

Language

English

Abstract

A number of nitrogen-containing fused heterocyclic compds., such as triazolotriazine, triazinotriazine, imidazotriazine, triazinoquinazolinone, and 1,2,4-triazine derivs. have been synthesized from 5-benzyl-6-chloro-3-(4-methylphenyl)-1,2,4-triazines. The antimicrobial activity was determined for thirteen representative compds.

L13 ANSWER 2 OF 39 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Comparative study on the reactivity of 6-haloimidazo[1,2-a]pyridine derivatives towards Negishi- and Stille-coupling reactions

Author/Inventor

Hervet, Maud; Thery, Isabelle; Gueffier, Alain; Enguehard-gueffier, Cecile
Patent Assignee/Corporate Source
Laboratoire de Chimie Therapeutique EA 3247, Faculte de Pharmacie, Tours, F-37200, Fr.

Source
Helvetica Chimica Acta (2003), 86(10), 3461-3469 CODEN: HCACAV; ISSN: 0018-019X

Document Type
Journal

Language
English

Abstract
The scope of the Suzuki-cross-coupling reaction of 6-haloimidazo[1,2-a]pyridines is dependent on the availability of the (hetero)arylboronic acids. Thus, with the aim to develop expanded applications of (hetero)arylations of imidazo[1,2-a]pyridines, we investigated the Negishi- and Stille-cross-coupling reactions at the 6-position. Remarkably, attempts to apply the Negishi-cross-coupling conditions to the organozinc derivative prepared from 6-haloimidazo[1,2-a]pyridine via a lithium-zinc exchange led to the 5-Ph compound 3 in 54% yield instead of the desired 6-Ph isomer (Scheme 1). In contrast, various com. available halogenated five- or six-membered-ring heterocycles were efficiently coupled to the 6-(trialkylstannyl)imidazo[1,2-a]pyridine under Stille conditions (Table 2).

L13 ANSWER 3 OF 39 CAPLUS COPYRIGHT 2008 ACS on STN
Title

Ips- or Cine-Substitutions of 6-Haloimidazo[1,2-a]pyridine Derivatives with Different Azoles Depending on the Reaction Conditions

Author/Inventor
Enguehard, Cecile; Allouchi, Hassan; Gueffier, Alain; Buchwald, Stephen L.

Patent Assignee/Corporate Source
Laboratoire de Chimie Therapeutique et Laboratoire de Chimie Physique, UFR des Sciences Pharmaceutiques, Tours, 37200, Fr.

Source
Journal of Organic Chemistry (2003), 68(14), 5614-5617 CODEN: JOCEAH; ISSN: 0022-3263

Document Type
Journal

Language
English

Abstract
The reactivity of 6-haloimidazo[1,2-a]pyridines I (R = Br, iodo) toward different azoles, such as pyrrole, indole, 1,2,3-triazole, etc., is reported. The process was shown to be highly dependent on the reaction conditions. Thus, in the presence of a copper(I) catalyst, the product of ipso substitution was obtained, whereas in the absence of copper, with cesium carbonate in N,N-dimethylformamide, a cine substitution took place.

L13 ANSWER 4 OF 39 CAPLUS COPYRIGHT 2008 ACS on STN
Title

Easy Access to Novel Substituted 6-Aminoimidazo[1,2-a]pyridines Using Palladium- and Copper-Catalyzed Aminations

Author/Inventor
Enguehard, Cecile; Allouchi, Hassan; Gueffier, Alain; Buchwald, Stephen L.

Patent Assignee/Corporate Source
EA 3247, Laboratoire de Chimie Therapeutique, Laboratoire de Chimie Physique, UFR des Sciences Pharmaceutiques, Tours, 37200, Fr.

Source
Journal of Organic Chemistry (2003), 68(11), 4367-4370 CODEN: JOCEAH; ISSN: 0022-3263

Document Type
Journal

Language
English

Abstract
Novel 6-aminoimidazo[1,2-a]pyridines I (R1 = Ph, PhCH2, n-hexyl, cyclohexyl, R2 = H; R1 = Ph, R2 = Me; R1R2N = pyrrolidino, piperidino, morpholino, N-Et piperazino) were readily prepared via palladium- or copper-catalyzed amination of the corresponding 6-haloimidazo[1,2-a]pyridines with primary and secondary amines R1NR2.

L13 ANSWER 5 OF 39 CAPLUS COPYRIGHT 2008 ACS on STN
Title

Preparation of imidazo[1,2-a]pyridines for the prophylaxis and treatment of herpes viral infections

Author/Inventor
Gudmundsson, Kristjan; Johns, Brian A.

Patent Assignee/Corporate Source
Smithkline Beecham Corporation, USA

Source
PCT Int. Appl., 144 pp. CODEN: PIXXD2

Document Type
Patent

Language
English

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
------------	------	------	-----------------	------

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003000689	A1	20030103	WO 2002-US18520	20020610

Patent Number (1)
WO 2003000689
Kind Code (1)
A1
Patent Publication Date (1)
20030103
Application Number (1)
WO 2002-US18520
Application Date (1)
20020610
Priority Patent Number (1)
US 2001-300009P
Priority Kind Code (1)
P
Priority Patent Publication Date (1)
20010621

Abstract

The title compds. [I; p = 0-4; R1 = halo, alkyl, alkenyl, etc.; R2 = halo, alkenyl, cycloalkyl, etc.; Y = N, CH; R3, R4 = H, halo, alkyl, etc.; q = 0-5; R5 = halo, alkyl, alkenyl, etc.] were prepared. E.g., a 7-step synthesis of II, starting from 2-amino-3-nitropyridine and 2-bromo-4'-fluoroacetophenone, which showed IC50 of 0.6 μ M against HSV-1, was given.

L13 ANSWER 6 OF 39 CAPLUS COPYRIGHT 2008 ACS on STN

Title
Product class 5: azaindolizines with two nitrogen atoms in the five-membered ring
Author/Inventor
Hajos, G.; Riedl, Z.
Patent Assignee/Corporate Source
Chemical Research Center, Institute of Chemistry, Budapest, H-1025, Hung.
Source
Science of Synthesis (2002), 12, 613-678 CODEN: SSCYJ9
Document Type
Journal; General Review
Language
English

Abstract

A review of preparation of azaindolizines with two nitrogen atoms in the five-membered ring. Covered reactions include ring-closure, substituent modification, substitution reactions, and other miscellaneous methods.

L13 ANSWER 7 OF 39 CAPLUS COPYRIGHT 2008 ACS on STN

Title
Preparation of imidazolyl derivatives as agonists or antagonists of somatostatin receptors
Author/Inventor
Thurieu, Christophe Alain; Poitout, Lydie Francine; Galcera, Marie-Odile; Gordon, Thomas D.; Morgan, Barry A.; Moinet, Christophe Philippe; Bigg, Dennis
Patent Assignee/Corporate Source
Societe De Conseils De Recherches Et D'applications Scientifiques (S.C.R.A.S.), Fr.
Source
PCT Int. Appl., 369 pp. CODEN: PIXXD2
Document Type
Patent
Language
English

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002010140	A2	20020207	WO 2001-US23959	20010731

Patent Number (1)
WO 2002010140
Kind Code (1)
A2
Patent Publication Date (1)
20020207
Application Number (1)
WO 2001-US23959
Application Date (1)
20010731
Priority Patent Number (1)
US 2000-222584P
Priority Kind Code (1)

P
Priority Patent Publication Date (1)
20000801

Abstract Imidazole derivs. I [R1 = H, (CH2)mCO(CH2)mZ1, (CH2)mZ1, etc.; Z1 = (un)substituted benzo[b]thiophene, Ph, naphthyl, etc.; m = 0-6; R2 = H, alkyl; R1 and R2 taken together with the nitrogen atoms to which they are attached form II-IV; R3 = (CH2)mE(CH2)mZ2; E = O, S, CO, etc.; Z2 = H, alkyl, NH2, etc.; R4 = H, (CH2)mA1; A1 = C(Y)NX1X2; C(NH)X2, X2; Y = O, S; X1 = H, alkyl, etc.; X2 = alkyl, etc.; R5 = alkyl, (un)substituted aryl, etc.; R6 = H, alkyl; R7 = alkyl, (CH2)mZ4; Z4 = (un)substituted Ph, naphthyl, indolyl, etc.], which are useful as agonists or antagonists of somatostatin receptors (no data) and for inhibiting the proliferation of *Helicobacter pylori*, were prepared. Thus, activating 2-furancarboxylic acid with carbonyldiimidazole followed by addition of 2-[(1S)-1-amino-2-(indol-3-yl)ethyl]-4-phenyl-1H-imidazole afforded 94% the title compound V. Compds. I are effective at 0.01-10.0 mg/kg/day.

L13 ANSWER 8 OF 39 CAPLUS COPYRIGHT 2008 ACS on STN

Title (Hetero)arylation of 6-halogenoimidazo[1,2-a]pyridines differently substituted at C(2): influence of the 2-substituent on the Suzuki cross-coupling reaction

Author/Inventor Enguehard, Cecile; Hervet, Maud; Thery, Isabelle; Renou, Jean-Louis; Fauvel, Florence; Gueiffier, Alain
Patent Assignee/Corporate Source EA 3247, Laboratoire de Chimie Therapeutique, UFR des Sciences Pharmaceutiques, Tours, F-37200, Fr.

Source Helvetica Chimica Acta (2001), 84(12), 3610-3615 CODEN: HCACAV; ISSN: 0018-019X

Document Type Journal

Language English

Abstract The authors previously reported that reactivity towards the Suzuki cross-coupling reaction of 3-iodoimidazo[1,2-a]pyridines substituted at C(2) is largely influenced by the nature of this 2-substituent. With the aim to expand the scope of this coupling process to the 6-position of this series, it seemed important to similarly determine the influence of the nature of the 2-substituent (H, alkyl, or aryl) on the rate of coupling. From this work, the Suzuki-type cross-coupling proceeds efficiently on 6-bromo-2-methyl- and 2-(4-fluorophenyl)imidazo[1,2-a]pyridines, whereas the 6-Br derivative unsubstituted at C(2) appeared to be poorly reactive. By modifying the reaction conditions in terms of catalyst and base, and the nature of the halogen, the reactivity of the unsubstituted series was largely enhanced. This work established efficient and convenient Suzuki reaction conditions for the 6-(hetero)arylation of 6-haloimidazo[1,2-a]pyridines depending on the nature of the 2-substituent and boronic acid.

L13 ANSWER 9 OF 39 CAPLUS COPYRIGHT 2008 ACS on STN

Title Reactivity of a 6-chloroimidazo[1,2-b]pyridazine derivative towards Suzuki cross-coupling reaction

Author/Inventor Enguehard, Cecile; Hervet, Maud; Allouchi, Hassan; Debouzy, Jean-Claude; Leger, Jean-Michel; Gueiffier, Alain
Patent Assignee/Corporate Source Laboratoire de Chimie Therapeutique, Faculte de Pharmacie, Tours, 37200, Fr.

Source Synthesis (2001), (4), 595-600 CODEN: SYNTBF; ISSN: 0039-7881

Document Type Journal

Language English

Abstract The influence of base, reaction time and boronic acid in the Suzuki cross-coupling reaction on 6-chloro-2-(4-fluorophenyl)imidazo[1,2-b]pyridazine is reported. The crystal data for the thien-2-yl compound I is also described.

L13 ANSWER 10 OF 39 CAPLUS COPYRIGHT 2008 ACS on STN

Title Synthesis of substituted imidazopyrazines as ligands for the human somatostatin receptor subtype 5

Author/Inventor Contour-Galceran, M.-O.; Poitout, L.; Molinet, C.; Morgan, B.; Gordon, T.; Roubert, P.; Thureau, C.
Patent Assignee/Corporate Source Institut Henri Beaufour, Les Ulis, F-91966, Fr.

Source Bioorganic & Medicinal Chemistry Letters (2001), 11(5), 741-745 CODEN: BMCL8; ISSN: 0960-894X

Document Type Journal

Language English

Abstract A new preparation of trisubstituted imidazopyrazines and dihydroimidazopyrazines via parallel synthesis using amino acids and bromo ketones resulted in the discovery of non-peptidic sst5 selective agonists.

L13 ANSWER 11 OF 39 CAPLUS COPYRIGHT 2008 ACS on STN

Title Preparation of imidazoline derivatives for the treatment of diabetes, especially type II diabetes
 Author/Inventor Paal, Michael; Ruehter, Gerd; Schotten, Theo
 Patent Assignee/Corporate Source Eli Lilly and Company, USA
 Source PCT Int. Appl., 143 pp. CODEN: PIXXD2
 Document Type Patent
 Language English

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000078726	A1	20001228	WO 2000-US11881	20000619

Patent Number (1)

WO 2000078726

Kind Code (1)

A1

Patent Publication Date (1)

20001228

Application Number (1)

WO 2000-US11881

Application Date (1)

20000619

Priority Patent Number (1)

GB 1999-14222

Priority Kind Code (1)

A

Priority Patent Publication Date (1)

19990618

Abstract

The title compds. [I; R1-R4 = H, alkyl; R1 and R3, together with the carbon atoms to which they are attached, combine to form a C3-7 carbocyclic ring and R2 and R4 = H, alkyl; R1 and R2, together with the carbon atom to which they are attached combine to form a C3-7 spirocarbocyclic ring and R3 and R4 = H, alkyl; R3 and R4, together with the carbon atom to which they are attached combine to form a C3-7 spirocarbocyclic ring and R1 and R2 = H, alkyl; R5 = H, alkyl, aryl, etc.; R6 = H, alkyl, alkoxy, etc.; R7 = H, alkyl, alkoxy, etc.; Y = NHCONH, NHCO, a bond, etc.; A = a monocyclic or bicyclic ring; R8 = H, alkyl, alkenyl, etc.; R9, R10 = H, alkyl, alkoxy, etc.], useful for the treatment of diabetes, diabetic complications, metabolic disorders, or related diseases where impaired glucose disposal is present (no data), were prepared and formulated. E.g., a multi-step synthesis of the imidazoline II.HCl was given. The compds. I are effective at 0.1-5 mg/kg/day.

L13 ANSWER 12 OF 39 CAPLUS COPYRIGHT 2008 ACS on STN

Title Preparation of imidazolyl derivatives as agonists or antagonists of somatostatin receptors
 Author/Inventor Thirieau, Christophe Alain; Poltout, Lydie Francine; Galcera, Marie-Odile; Gordon, Thomas D.; Morgan, Barry; Moinet, Christophe Philippe
 Patent Assignee/Corporate Source Societe de Conseils de Recherches et d'Applications Scientifiques, S.A., Fr.
 Source PCT Int. Appl., 342 pp. CODEN: PIXXD2
 Document Type Patent
 Language English

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9964401	A2	19991216	WO 1999-US12760	19990608

Patent Number (1)

WO 9964401

Kind Code (1)

A2

Patent Publication Date (1)

19991216

Application Number (1)

WO 1999-US12760

Application Date (1)

19990608

Priority Patent Number (1)

US 1998-89087P
Priority Kind Code (1)

P
Priority Patent Publication Date (1)
19980612

Abstract

The title compds. [I: R1 = H, (CH₂)mCO(CH₂)mZ1, (CH₂)mZ1, etc.; Z1 = (un)substituted benzo[b]thiophene, Ph, naphthyl, etc.; R2 = H, alkyl; R1 and R2 taken together with the nitrogen atoms to which they are attached form II-IV; R3 = (CH₂)mE(CH₂)mZ2; E = O, S, CO, etc.; Z2 = H, alkyl, NH₂, etc.; R4 = H, (CH₂)mA1; A1 = C-(Y)NX1X2; C-(Y)X2; C-(NH)X2, X2; Y = O, S; X1 = H, alkyl, etc.; X2 = alkyl, etc.; R5 = alkyl, (un)substituted aryl, etc.; R6 = H, alkyl; R7 = alkyl, (CH₂)mZ4; Z4 = (un)substituted Ph, naphthyl, indolyl, etc.; m = 0-6] which are useful as agonists or antagonists of somatostatin receptors (no data), and for inhibiting the proliferation of *Helicobacter pylori*, were prepared. Thus, activating 2-furancarboxylic acid with carbonyldimidazole followed by addition of 2-(1S)-1-amino-2-(indol-3-yl)ethyl]-4-phenyl-1H-imidazole afforded 94% the title compound V. Compds. I are effective at 0.01-10.0 mg/kg/day.

L13 ANSWER 13 OF 39 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Imidazo[1,2-b]pyridazines: syntheses and interaction with central and peripheral-type (mitochondrial) benzodiazepine receptors

Author/Inventor

Barlin, Gordon B.

Patent Assignee/Corporate Source

Division of Neuroscience, John Curtin School of Medical Research, Australian National University, Canberra, ACT 2601, Australia

Source

Journal of Heterocyclic Chemistry (1998), 35(5), 1205-1217 CODEN: JHTCAD; ISSN: 0022-152X

Document Type

Journal

Language

English

Abstract

The fundamental chemical of pyridazines, the syntheses of substituted imidazo[1,2-b]pyridazines (1) (and some related compds.) and the interaction of the products with central benzodiazepine receptors (CBR) and peripheral-type (mitochondrial) benzodiazepine receptors (PBR) are described. Some of these imidazo[1,2-b]pyridazines had high selective affinity for the central benzodiazepine receptors and others had high selectivity for the peripheral-type (mitochondrial) benzodiazepine receptors. The results of structure-activity studies and mol. modeling will be reported. In vivo tests of some compds. which interacted strongly with the central benzodiazepine receptors revealed reasonably potent anticonvulsant/anticonflict activity, and some of those which bind selectively to the peripheral-type (mitochondrial) benzodiazepine receptors are being examined as possible radiopharmaceuticals for imaging of tumors (and other disease states).

L13 ANSWER 14 OF 39 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Bis-Cationic heteroaromatics as macrofilaricides: synthesis of bis-amidine and bis-guanyldiazene derivatives of substituted imidazo[1,2-a]pyridines

Author/Inventor

Sundberg, Richard J.; Biswas, Sujay; Murthi, Krishna Kumar; Rowe, Donna; McCall, John W.; Dzimiński, Michael T.

Patent Assignee/Corporate Source

Department of Chemistry, University of Virginia, Charlottesville, VA, 22901, USA

Source

Journal of Medicinal Chemistry (1998), 41(22), 4317-4328 CODEN: JMCMAR; ISSN: 0022-2623

Document Type

Journal

Language

English

Abstract

A series of guanyldiazene, amidine, and hydrazone derivs. of 2-phenylimidazo[1,2-a]pyridine have been prepared and evaluated for macrofilarial activity against *Acanthocheilonema viteae* and *Brugia pahangi* in jirds. Compds. with 4',6-bis-substitution by cyclic guanyldiazene groups show activity. 4',6-Bis-amidines show some activity but are more toxic; 4' or 6-monosubstituted compds. are inactive. 2,6-Bis-substituted compds. lacking the Ph ring are inactive. 4',6-Bis-substituted compds. having addnl. double bonds inserted between the heterocyclic ring and the Ph ring or between the substituent and the ring system show reduced activity.

L13 ANSWER 15 OF 39 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Preparation of fused imidazopyridines as antiulcer drugs

Author/Inventor

Tanaka, Hironori; Fukuzumi, Kazuyoshi; Togawa, Takeshi; Banno, Kimiko; Ushiro, Toshihisa; Morii, Masaaki; Nakatani, Takafumi

Patent Assignee/Corporate Source

Shinippon Pharmaceutical Inc., Japan

Source

PCT Int. Appl., 145 pp. CODEN: PIXXD2

Document Type

Patent

Language

Japanese

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9633195	A1	19961024	WO 1996-JP975	19960410

Patent Number (1)

WO 9633195

Kind Code (1)

A1

Patent Publication Date (1)

19961024

Application Number (1)

WO 1996-JP975

Application Date (1)

19960410

Priority Patent Number (1)

JP 1995-97130

Priority Kind Code (1)

A

Priority Patent Publication Date (1)

19950421

Abstract

The title compds. [I: A or B ring = benzene, thiophene, furan, or pyrrole ring; R1 = OH, halo, lower (halo)alkyl, lower alkoxy, acyloxy; k = 0-3; R2, R3 = H, alkenyl, acyl, alkoxycarbonyl, (un)substituted amino, aryl (un)substituted alkyl; or R2 and R3 combine with N to form a N-containing heterocycle, etc.; R4, R5 = halo, cyano, OH, CO2H, acyl, etc.; m = 0-2; n = 0-2; broken line may combine with solid line to represent a single or double bond], pharmacol. acceptable salts thereof, and solvates thereof are prepared. I are useful as anticancer agents and for treating gastrointestinal diseases. 2-(2-Methylphenyl)imidazo[1,2-a]quinoline was treated with NaNO2 and then with Zn to give I [A(R1)k = 2-MeC6H4, NR2R3 = NH2, R4 = R5 = H, broken line combine with solid line to represent a double bond] (II). II showed IC50 of 26.0 µM against H⁺/K⁺-ATPase and inhibited stomach acid secretion in rats and ulcer formation in mice.

L13 ANSWER 16 OF 39 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Imidazo[1,2-b]pyridazines. XIX. Syntheses and central nervous system activities of some 6-arythio(aryloxy and alkylthio)-3-(acetamidomethyl, benzamidomethyl, methoxy and unsubstituted)-2-arylimidazo[1,2-b]pyridazines

Author/Inventor

Barlin, Gordon B.; Davies, Les P.; Ireland, Stephen J.

Patent Assignee/Corporate Source

Div. Neuroscience, Australian National Univ., Canberra, 2601, Australia

Source

Australian Journal of Chemistry (1996), 49(4), 443-449 CODEN: AJCHAS; ISSN: 0004-9425

Document Type

Journal

Language

English

Abstract

Some 6-arythio(aryloxy and alkylthio)-3-(acetamidomethyl, benzamidomethyl, methoxy and unsubstituted)-2-arylimidazo[1,2-b]pyridazines have been prepared and examined for their ability to displace [3H]diazepam from rat brain membranes. The most active compound was 3-acetamidomethyl-2-(3',4'-methylenedioxyphenyl)-6-phenylthioimidazo[1,2-b]pyridazine with IC50 4.4 nM. The 3-acylaminoethyl-6-(2- and 3-methoxyphenylthio)-2-phenylimidazo[1,2-b]pyridazines proved less active than their 6-phenylthio analogs, and larger substituents at the 2- and 6-positions markedly decreased binding. Significant differences in binding ability have been observed between 3-acylaminoethyl-2-aryl-6-phenylthioimidazo[1,2-b]pyridazines and the corresponding imidazo[1,2-a]pyridines.

L13 ANSWER 17 OF 39 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Sterically crowded heterocycles. IV. Diastereoisomeric products by ferricyanide oxidation of quaternary pyridinium salts

Author/Inventor

Kubik, Richard; Kuthan, Josef

Patent Assignee/Corporate Source

Dep. Org. Chem., Prague Inst. Chem. Technol., Prague, 166 28, Czech Rep.

Source

Collection of Czechoslovak Chemical Communications (1996), 61(4), 615-621 CODEN: CCCGAK; ISSN: 0010-0765

Document Type

Journal

Language

English

Abstract

While ferricyanide oxidation of achiral 4-(4-dimethylaminophenyl)-2,6-diphenyl-1-(pyridin-2-yl)pyridinium perchlorate gave racemic 3-(4-dimethylaminophenyl)-1-phenyl-3-(2-phenylimidazo[1,2-a]pyridin-3-yl)prop-2-1-one, the same oxidative procedure applied to racemic 1-[5-(1-methylpyrrolidin-2-yl)pyridin-2-yl]-2,4,6-triphenylpyridinium perchlorate or its 4-(dimethylamino)phenyl derivative led to mixts. of diastereoisomeric 3-[6-(1-methylpyrrolidin-2-yl)-2-phenylimidazo[1,2-a]pyridin-3-yl]-1,3-diphenyl-2-en-1-ones I (X = H) or

their dimethylamino derivs. I (X = NMe₂), resp.

L13 ANSWER 18 OF 39 CAPLUS COPYRIGHT 2008 ACS on STN

Title Imidazo[1,2-b]pyridazines: Studies on chemical structure-antiinflammatory activity relationships
Author/Inventor Abignente, Enrico; Arena, Francesca; Luraschi, Elena; Sacchi, Antonia; Rimoli, Maria Grazia; Laneri, Sonia
Patent Assignee/Corporate Source Facolta di Farmacia, Universita di Napoli "Federico II", Naples, 80131, Italy
Source Acta Chimica Slovenica (1994), 41(2), 131-48 CODEN: ACSLE7; ISSN: 1318-0207
Document Type Journal
Language English

Abstract The synthesis and pharmacol. testing of some series of imidazo[1,2-b]pyridazine derivs. bearing an acidic function are described. The exptl. results are discussed, taking into particular consideration the pharmacol. profile of these new compds., which is chiefly characterized by the prevalence of the analgesic activity over the antiinflammatory activity. The relations between such pharmacol. activities and chemical structures are discussed.

L13 ANSWER 19 OF 39 CAPLUS COPYRIGHT 2008 ACS on STN

Title Reaction of 1-aminophthalazines with α -halocarbonyl compounds: Imidazo-2,1-a]phthalazines and their benzodiazepine receptor activities
Author/Inventor Catarzi, Daniela; Cecchi, Lucia; Colotta, Vittoria; Conti, Gabriele; Melani, Fabrizio; Filacchioni, Guido; Martini, Claudia; Giusti, Laura; Lucacchini, Antonio
Patent Assignee/Corporate Source Dip. Sci. Farm., Univ. Firenze, Florence, 50121, Italy
Source Farmaco (1993), 48(4), 447-57 CODEN: FRMCE8; ISSN: 0014-827X
Document Type Journal
Language English

Abstract The one-pot synthesis and the benzodiazepine receptor binding activity of some imidazo[2,1-a]phthalazines (I, R = H, Me, Ph, or NH₂, R₁ = H, p-MeOC₆H₄, CO₂Et, etc., R₂ = H, Ph or CO₂Et) bearing different substituents at position-2 and/or 3 and/or 6 is reported. The dissimilar binding results of the reported compds. are discussed in relation to the nature and/or position of the substituents on the tricyclic ring system.

L13 ANSWER 20 OF 39 CAPLUS COPYRIGHT 2008 ACS on STN

Title Imidazo[1,2-a]pyridines. I. Synthesis and inotropic activity of new 5-imidazo[1,2-a]pyridinyl-2(1H)-pyridone derivatives. [Erratum to document cited in CA115(15):159040c]
Author/Inventor Yamanaka, Motosuke; Miyake, Kazutoshi; Suda, Shinji; Ohhara, Hideto; Ogawa, Toshiaki
Patent Assignee/Corporate Source Eisai Tsukuba Res. Lab., Tsukuba, 300-26, Japan
Source Chemical & Pharmaceutical Bulletin (1991), 39(12), 3382 CODEN: CPBTAL; ISSN: 0009-2363
Document Type Journal
Language English

Abstract An error in Chart 3 has been corrected. An error in Table III has been corrected. Three errors in the text have been corrected. The errors were not reflected in the abstract but were reflected in the index entries.

L13 ANSWER 21 OF 39 CAPLUS COPYRIGHT 2008 ACS on STN

Title Imidazo[1,2-a]pyridines. I. Synthesis and inotropic activity of new 5-imidazo[1,2-a]pyridinyl-2(1H)-pyridone derivatives
Author/Inventor Yamanaka, Motosuke; Miyake, Kazutoshi; Suda, Shinji; Ohhara, Hideto; Ogawa, Toshiaki
Patent Assignee/Corporate Source Eisai Tsukuba Res. Lab., Tsukuba, 300-26, Japan
Source Chemical & Pharmaceutical Bulletin (1991), 39(6), 1556-67 CODEN: CPBTAL; ISSN: 0009-2363
Document Type

Journal
Language English
Abstract

A series of imidazo[1,2-a]pyridinylpyridinones, e.g., I (R1 = H, Br, Cl cyano; R2 = Me, Et; R3 = H, 6-Cl, 6-F, 6-cyano, 6-Me, 6-OMe, 6-CF3, 8-F) and II (R4 = H, Me, CH2OMe, Ph; R5 = H, Me, R6 = Me, Et) was synthesized and evaluated for pos. inotropic activity. II (R4 = R5 = H, R6 = Me) (III) was a potent and selective inhibitor of phosphodiesterase III and a long-acting, potent, orally pos. inotropic agent. Addnl. imidazo[1,2-a]pyridin-2-yl compds. were also prepared. Altering the pyridine substitution from I to II produced a 2-fold increase in the i.v. cardiotoxic potency (ED50) from 52 to 23 µg/kg, while substitution at the 3-, 7- or 8-position reduced potency. In I, introduction of halogen groups enhanced the activity and I (R1 = Cl, R2 = Me, R3 = 6-F) was the most potent (i.v. ED50 11 µg/kg) in this series. III is presently under development for the treatment of congestive heart failure.

L13 ANSWER 22 OF 39 CAPLUS COPYRIGHT 2008 ACS on STN

Title Research on heterocyclic compounds. XXVI. Anti-inflammatory and related activities of some 2-phenylimidazo[1,2-b]pyridazines
Author/Inventor Abignente, Enrico; Arena, Francesca; Luraschi, Elena; De Caprariis, Paolo; Marmo, Emilio; Vitagliano, Saverio; Donnoli, Donato
Patent Assignee/Corporate Source Fac. Farm., Univ. Napoli "Federico II", Naples, 80131, Italy
Source Research Communications in Chemical Pathology and Pharmacology (1990), 67(1), 43-54 CODEN: RCOCB8; ISSN: 0034-5164
Document Type Journal
Language English
Abstract

Five acidic Ph derivs. of the imidazo[1,2-b]pyridazines I (R and R1 = H or Me, R2 = H, Ph or C6H4OMe-4) system were subjected to some tests in vivo in order to evaluate their biol. activity. Anti-inflammatory activity was studied by means of the carrageenin rat paw edema, whereas writhing induced in mice by acetic acid was used to assess analgesic activity. The irritative and ulcerogenic action on the rat gastric mucosa was examined after oral administration of larger doses. The inhibitory activity on platelet malondialdehyde production was studied in vitro. The exptl. results are discussed from the point of view of structure-activity relationships and mode of action.

L13 ANSWER 23 OF 39 CAPLUS COPYRIGHT 2008 ACS on STN

Title Synthesis and cardiotonic activity of 2,4-diamino-1,3,5-triazines
Author/Inventor Kosary, Judit; Kasztreiner, Endre; Rabclozky, Gyorgy; Kurthy, Maria
Patent Assignee/Corporate Source Inst. Drug Res., Budapest, H-1325, Hung.
Source European Journal of Medicinal Chemistry (1989), 24(1), 97-9 CODEN: EJMCA5; ISSN: 0223-5234
Document Type Journal
Language English
Abstract

Thirty-two diamino-1,3,5-triazine derivs., e.g., I (R = NH2, NHBu, NMe2, morpholino, 4-methylpiperazino; R1 = indol-3-yl, imidazo[1,2-a]pyridin-2-yl, imidazo[1,2-b]pyridazin-2-yl, imidazo[1,2-a]pyrazin-2-yl, etc.) were prepared and tested for cardiotonic activity in cats. There was no correlation between the cardiotonic activity of these compds. and their inhibitory effect on cardiac phosphodiesterase.

L13 ANSWER 24 OF 39 CAPLUS COPYRIGHT 2008 ACS on STN

Title Preparation of imidazo[1,2-b]pyridazines as nervous system agents
Author/Inventor Barlin, Gordon Bruce; Davies, Leslie Philip; Ireland, Stephen James; Ngu, Maria Mee Leng
Patent Assignee/Corporate Source Australian National University, Australia
Source PCT Int. Appl., 66 pp. CODEN: PIXXD2
Document Type Patent
Language English
Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 8901333	A1	19890223	WO 1988-AU290	19880805

Patent Number (1)
WO 8901333
Kind Code (1)

A1
 Patent Publication Date (1)
 19890223
 Application Number (1)
 WO 1988-AU290
 Application Date (1)
 19880805
 Priority Patent Number (1)
 AU 1987-3617
 Priority Kind Code (1)
 A
 Priority Patent Publication Date (1)
 19870807

Abstract

The title compds. [I; X = halo, ER1; E = O, S, NH, OCH2, SCH2, NHCH2; R1 = (un)substituted alkyl, alkenyl, alkynyl, cycloalkyl, aryl, aralkyl, heterocyclyl; Y = OR2, CH2NHCOR3, CH2NR4R5; R2-R5 = alkyl, cycloalkyl, aryl; Z = substituted alkyl, (un)substituted cycloalkyl, aryl, aralkyl, heterocyclyl] were prepared as nervous system agents useful, e.g., as anxiolytics, muscle relaxants, anticonvulsants, and for treatment of insomnia and reversal of the sedative effects of benzodiazepines. A mixture of 0.5 g 6-fluoro-3-pyridazinamine, 0.74 g 4-MeC6H4COCHO.H2O, and 0.5 mL concentrated HCl was refluxed 5.5 h in EtOH to give I (X = F, Y = OH, Z = 4-MeC6H4) which was etherified with CH2N2 in Et2O to give 0.6 g II (X = F, Y = MeO, Z = 4-MeC6H4) (II). II displaced diazepam from rat brain synaptosomal membrane with an IC50 of 17 nM.

L13 ANSWER 25 OF 39 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Aryloxy- and aralkylthio-substituted imidazo [1,2-b]pyridazines with benzodiazepine receptor-binding activity, their preparation, pharmaceutical compositions, and use

Author/Inventor

Barlin, Gordon Bruce; Davies, Leslie Philip; Ireland, Stephen James; Ngu, Maria Mee Leng

Patent Assignee/Corporate Source

Australian National University, Australia; University of Sydney; Australasian Drug Development Ltd.

Source

PCT Int. Appl., 44 pp. CODEN: PIXXD2

Document Type

Patent

Language

English

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 8901478	A1	19890223	WO 1988-AU291	19880805

Patent Number (1)

WO 8901478

Kind Code (1)

A1

Patent Publication Date (1)

19890223

Application Number (1)

WO 1988-AU291

Application Date (1)

19880805

Priority Patent Number (1)

AU 1987-3617

Priority Kind Code (1)

A

Priority Patent Publication Date (1)

19870807

Abstract

Title compds. I [X = OAr, SCH2Ar; Ar = (un)substituted aryl; Y = alkoxy; Z = (un)substituted (hetero)aryl], which bind to diazepam receptors, were prepared. Treatment of 3-MeOC6H4CH2Cl with thiourea followed by basic hydrolysis gave 3-MeOC6H4CH2SH, which was condensed with 6-chloropyridazin-3-amine 2-oxide in 0.5M NaOH to give 6-(3-methoxybenzylthio)pyridazin-3-amine 2-oxide. Cyclocondensation of the latter with 3-(bromoacetyl)pyridine-HBr and methylation of the obtained alc. (Y = OH) with CH2N2 gave I (X = SCH2C6H4OMe-3, Y = OMe, Z = 3-pyridyl) (II). The IC50 of II for displacement of specifically bound [3H]-diazepam from rat brain benzodiazepine receptors in the presence of 100 µM GABA was 5 nM (cf. 4.2 nM for diazepam).

L13 ANSWER 26 OF 39 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Preparation of 1-[imidazo[1,2-a]pyridin-6-yl]-2-propanones as intermediates for cardiotonics

Author/Inventor

Yamanaka, Motosuke; Myake, Kazutoshi; Suda, Shinji; Ohara, Hideto; Ogawa, Toshiaki

Patent Assignee/Corporate Source

Eisai Co., Ltd., Japan

Source

Document Type
Patent
Language

Japanese

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 63077879	A	19880408	JP 1986-221885	19860922

Patent Number (1)
JP 63077879
Kind Code (1)
A
Patent Publication Date (1)
19880408
Application Number (1)
JP 1986-221885
Application Date (1)
19860922
Priority Patent Number (1)
JP 1986-221885
Priority Patent Publication Date (1)
19860922

Abstract

Title compds. I [R1 = H, alkyl, R5CH2 (R5 = alkyl), R6R7N (R6, R7 = H, alkyl), Ph; R2 = H, halo; R3 = H, alkyl, halo; R4 = H, alkyl), useful intermediates for cardiotonics II and III (X = H, Me, Y = cyano, carboxamide, H, Nt2, halo; Z = H, alkyl; W = H, alkyl), are prepared Ozonolysis of 6-(2-methyl-2-propenyl)imidazo[1,2-a]pyridine (preparation given) in concentrated HCl-H2O-EtOH, followed by treatment with Na2NO3 gave 70.5% 1-[imidazo[1,2-a]pyridine-6-yl]-2-propanone, which in DMF was heated with DMF di-Me acetal at 80° for 1 h to afford 74.5% 4-dimethylamino-3-[imidazo[1,2-a]pyridine-6-yl]-3-butene-2-one (IV). A mixture of IV in DMF was heated with NCH2CONH2 and MeONa at 80-90° for 12 h to give III (R1 = R2 = R3 = Z = H; Y = CN; W = Me), which at 100 µg/kg showed 97%, 22%, and -17% changes for cardiovascular contraction, heart beat, and blood pressure in dogs.

L13 ANSWER 27 OF 39 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Imidazo[1,2-b]pyridazines. II. 6-(Alkylthio)- and 6-(arylthio)-3-methoxy-2-phenylimidazo[1,2-b]pyridazines

Author/Inventor

Barlin, Gordon B.; Ireland, Stephen J.

Patent Assignee/Corporate Source

John Curtin Sch. Med. Res., Aust. Natl. Univ., Canberra, 2601, Australia

Source

Australian Journal of Chemistry (1987), 40(8), 1491-7 CODEN: AJCHAS; ISSN: 0004-9425

Document Type

Journal

Language

English

Abstract

A series of 6-(alkylthio)- and 6-(arylthio)-3-methoxy-2-phenylimidazo[1,2-b]pyridazines I (e.g., R = Me, Et, Ph, p-ClC6H4, p-Me2NC6H4) were prepared from the corresponding 3-aminopyridazines II by sequential cyclocondensation with PhCOCHO and methylation with CH2N2.

L13 ANSWER 28 OF 39 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Preparation of imidazopyridazinylacrylamides as antihypertensives, diuretics, and saluretics

Author/Inventor

Meyer, Horst; Ingendoh, Axel; Garthoff, Bernhard; Hirth, Claudia

Patent Assignee/Corporate Source

Bayer A.-G., Fed. Rep. Ger.

Source

Ger. Offen., 22 pp. CODEN: GWXXBX

Document Type

Patent

Language

German

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 3542661	A1	19870604	DE 1985-3542661	19851203

Patent Number (1)
DE 3542661
Kind Code (1)
A1
Patent Publication Date (1)

19870604
 Application Number (1)
 DE 1985-3542661
 Application Date (1)
 19851203
 Priority Patent Number (1)
 DE 1985-3542661
 Priority Kind Code (1)
 A
 Priority Patent Publication Date (1)
 19851203

Abstract

The title compds. [I; R1 = H, alkoxy, (substituted) alkyl, alkenyl, aryl, heterocyclyl, alkylthio, sulfinyl, sulfonyl, amino, etc.; R2, R3 = H, alkyl; R4 = (substituted) aryl, heterocyclyl; R5 = H, alkyl, CF3; R6 = H, cyano, halo, alkoxy, alkyl, alkylsulfonyl, etc.; R7, R8 = H, aryl, (substituted) hydrocarbyl; R7R8N = heterocyclyl] were prepd as antihypertensives, diuretics, and saluretics. 3-Formyl-2-phenyl-6-piperidinoimidazo[1,2-b]pyridazine was added to a mixture of diethyl phosphonodiethylacetamide and NaH in C6H6 and the mixture was stirred 1 h at 45° to give 69% I (R1 = 1-piperidyl, R2 = R3 = R5 = R6 = H, R4 = Ph, NR7R8 = 1-piperidyl) had a urine output of 25 mL/kg containing 2900 µmol Na+/kg and 900 µmol K+/kg over 6 h.

L13 ANSWER 29 OF 39 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Imidazo[1,2-b]pyridazines. I. Some 3-alkoxy-6-halo-2-phenyl-(and 4'-substituted phenyl)imidazo[1,2-b]pyridazines and 3-methoxy-2,6-diphenylimidazo[1,2-b]pyridazine

Author/Inventor

Barlin, Gordon B.
 Patent Assignee/Corporate Source
 John Curtin Sch. Med. Res., Aust. Natl. Univ., Canberra, 2601, Australia

Source

Australian Journal of Chemistry (1986), 39(11), 1803-9 CODEN: AJCHAS; ISSN: 0004-9425

Document Type

Journal

Language

English

Abstract

A series of the title imidazopyridazines I (R = F, Cl, Br, Ph; R1 = Me, Et; R2 = H, Cl, Br, MeO; R3 = R4 = H) have been prepared from the relevant pyridazin-3-amines and arylglyoxals, followed by O-alkylation of the corresponding imidazo[1,2-b]pyridazin-3(5H)-ones with diazoalkanes. 6-Chloro-3-methoxy-2-phenylimidazo[2,1-a]phthalazine I (R = Cl; R1 = Me; R2 = H; R3R4 = CH:CHCH:CH) was prepared similarly.

L13 ANSWER 30 OF 39 CAPLUS COPYRIGHT 2008 ACS on STN

Title

3-Imidazo[1,2-a]pyridin-6-ylpyridine derivatives

Author/Inventor

Yamanaka, Motosuke; Miyake, Kazutoshi; Suda, Shinji; Ohara, Hideto; Ogawa, Toshiaki

Patent Assignee/Corporate Source

Eisai Co., Ltd., Japan

Source

Jpn. Kokai Tokkyo Koho, 21 pp. CODEN: JKXXAF

Document Type

Patent

Language

Japanese

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 61218589	A	19860929	JP 1985-59450	19850326

Patent Number (1)

JP 61218589

Kind Code (1)

A

Patent Publication Date (1)

19860929

Application Number (1)

JP 1985-59450

Application Date (1)

19850326

Priority Patent Number (1)

JP 1985-59450

Priority Kind Code (1)

A

Priority Patent Publication Date (1)

19850326

Abstract

The title compds. [I: X = H, Me; Y = cyano, CONH₂, H, NH₂, halo; W, Z = H, alkyl; R₁ = H, alkyl, alkoxyethyl, (un)substituted amino, etc.; R₂ = H, halo; R₃ = H, alkyl, halo], useful as cardiac stimulants, vasodilators, and blood platelet aggregation inhibitors, were prepared. Thus, a mixture of 4-(dimethylamino)-3-(imidazo[1,2-a]pyridin-6-yl)-3-buten-2-one, DMF, and NCCl₂CONH₂ was heated at 80-90° for 12 h to give 1,2-dihydro-5-(imidazo[1,2-a]pyridin-6-yl)-6-methyl-2-oxo-3-pyridinecarbonitrile. In vitro studies using guinea pig hearts showed that I at 10-4 to 10-6 M increased cardiac muscle contraction 34.1-110.1 %.

L13 ANSWER 31 OF 39 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Studies on heterocyclic compounds. XXII. The reaction of oxazolo[3,2-b]pyridazinium perchlorates with hydroxylamines

Author/Inventor

Sato, Kazue; Miyasaka, Tadashi

Patent Assignee/Corporate Source

Sch. Pharm. Sci., Showa Univ., Tokyo, 142, Japan

Source

Chemical & Pharmaceutical Bulletin (1983), 31(11), 3811-18 CODEN: CPBTAL; ISSN: 0009-2363

Document Type

Journal

Language

English

Abstract

The reaction of oxazolo[3,2-b]pyridazinium perchlorates (I: R = Me, Ph, OMe, Cl; R₁ = Ph, Me; R₂ = H, Me) with NH₂OH and KOH in DMF gave hydroxylaminopyridazines II (X = O, NOH), which were cyclized to imidazo[1,2-b]pyridazine 1-oxides III (X₁ = N=O) by heating in mineral acid. The N oxides were characterized by converting them into the corresponding imidazopyridazines III (X₁ = N) by deoxygenation.

L13 ANSWER 32 OF 39 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Synthesis of imidazo[1,2-b]pyridazine 1-oxides by the reaction of oxazolo[3,2-b]pyridazinium perchlorates with hydroxylamine

Author/Inventor

Sato, Kazue; Miyasaka, Tadashi; Arakawa, Kiichi

Patent Assignee/Corporate Source

Sch. Pharm. Sci., Showa Univ., Tokyo, Japan

Source

Chemistry Letters (1977), (12), 1501-4 CODEN: CMLTAG; ISSN: 0366-7022

Document Type

Journal

Language

English

Abstract

The reaction of oxazolo[3,2-b]pyridazinium perchlorates I (R = Ph, R₁ = Cl, Me, Ph; R = Me, R₁ = Ph) with HONH₂ gave monooximes II and dioximes III, from both of which were prepared imidazo[1,2-b]pyridazine 1-oxides IV (n = 1) by heating in mineral acid. The N-oxides were characterized by converting into IV (n = o) deoxygenation.

L13 ANSWER 33 OF 39 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Synthesis of 3-alkylthio-2-phenyl-5-(1-methyl-2-pyrrolyl)- and 3-alkylthio-2-phenyl-7-(1-methyl-2-pyrrolyl)pyrimidazoles

Author/Inventor

Godovikova, S. N.; Gol'dfarb, Ya. V.

Patent Assignee/Corporate Source

Inst. Org. Khim., Moscow, USSR

Source

Khim. Geterotsikl. Soedin., Sb. 1: Azotsoderzhashchie Geterotsikly (1967), 172-4. Editor(s): Hillers, S. Izd. "Zinatne": Riga, USSR. CODEN: 20NNA2

Document Type

Conference

Language

Russian

Abstract

The title compds. (I and II, resp.) were prepared for testing as fungicides by reaction of 3-bromo-2-phenyl-5-(1-methyl-2-pyrrolyl)pyrimidazole (III) and 3-bromo-2-phenyl-7-(1-methyl-2-pyrrolyl)pyrimidazole (IV) with BuLi, followed by reaction with alkyl disulfides. Thus, a mixture of 33.4 g, α-aminonitroline, 31.3 g, BrCH₂Br, and 200 ml, EtOH was refluxed 21 hrs. and worked up to give 66.4% 2-phenyl-5-(1-methyl-2-pyrrolyl)pyrimidazole (V), m. 112.5-13°; picrate m. 234-6°; HBr salt m. 258-60°. A solution of 15.6 g, Br in CHCl₃ was added to a solution of 27.1 g, V in CHCl₃, and the mixture stirred 2 hrs. and worked up to give 84.9% III, m. 74-6°; HBr salt m. 238-9°; picrate m. 191-3°. Similarly, 92.8% IV was prepared, m. 136.5-38°. To 10.7 g, III in 500 ml, Et₂O and 100 ml, C₆H₆ was added 0.033 mole BuLi in Et₂O, and the mixture refluxed 3 hrs., 0.033 mole alkyl disulfide in 20 ml, Et₂O added, and the mixture refluxed 9 hrs. and worked up to give the following I and II (compound type, R, m.p., % yield, m.p. picrate, and m.p. HBr salt given): I, Me, 91-2.5°, 61.4, 237-8°, 252°, I, Et, 108.5-9.2°, 90.5, 247°, 235-7°, I, Pr, -(oil), 74.8, 235-7°, 227-9°; II, Me, 94.5-5.5°, 62, 195-6°, 220-2°, and II, Et, -(oil), 55.6, 190-2°, 231-3°. Other compds. described are 5-bromo-2-phenyl-7-(1-methyl-2-pyrrolyl)pyrimidazole, m. 120-2° and an unknown compound, m. 141-2°, obtained in 80.5% yield from the reaction of III with 2

moles of BuLi, followed by reaction with Me2S.

L13 ANSWER 34 OF 39 CAPLUS COPYRIGHT 2008 ACS on STN

Title Imidazopyridazine derivatives
Author/Inventor Nitta, Yoshihiro; Yoneda, Fumio; Otaka, Takayuki
Patent Assignee/Corporate Source Chugai Pharmaceutical Co., Ltd.
Source 4 pp.
Document Type Patent
Language Unavailable

Patent Information				
PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 40022267	B4	19651002	JP	19631029

Patent Number (1)
JP 40022267
Kind Code (1)
B4
Patent Publication Date (1)
19651002
Application Number (1)
JP
Application Date (1)
19631029
Priority Patent Number (1)
JP
Priority Patent Publication Date (1)
19631029

Abstract

Manufacture of 6-R1-2-(R2-phenyl)imidazo[1,2-b]pyridazine (I), useful as inhibitor for central nerves, starting from the corresponding 6-halo compound was described. Thus, 2.3 g. 6-chloro-2-phenylimidazo[1,2-b]pyridazine is added to 100 ml. EtOH containing 0.9 g. NMe2, the whole heated at 160-70° for 3 hrs. in a sealed tube, an excess of EtOH removed, to the residue added H2O, and the precipitate recryst. from MeOH to give 2.1 g. I (R1 = NMe2, R2 = H), m. 195°. Similarly prepared are the following I (R1, R2, and m.p. given): NMe2, OMe, 204°; NMe2, Cl, 203°; NMe2, Br, 197°; morpholino, H, 183-4°; morpholino, OMe, 198-9°; morpholino, Cl, 227°; morpholino, Br, 243°; piperidino, H, 169°; piperidino, OMe, 153°; piperidino, Cl, 222°; piperidino, Br, 221°; pyrrolidino, H, 187-8°; pyrrolidino, OMe, 190°; pyrrolidino, Cl, 210°; pyrrolidino, Br, 211°.

L13 ANSWER 35 OF 39 CAPLUS COPYRIGHT 2008 ACS on STN

Title Synthesis of fused imidazo-heterocyclic systems
Author/Inventor Werbel, Leslie M.; Zamora, Maria L.
Patent Assignee/Corporate Source Parke, Davis & Co., Ann Arbor, MI
Source Journal of Heterocyclic Chemistry (1965), 2(3), 287-90 CODEN: JHTCAD; ISSN: 0022-152X
Document Type Journal
Language English

Abstract

The reaction of phenylacetyl bromide and a variety of α -amino- heterocycles was investigated to determine its applicability to the preparation of fused imidazo-heterocyclic systems. The imidazo[1,2-a]pyrazine, imidazo [1,2-b] pyridazine, and imidazo[1,2-a]-benzimidazole systems and some variations of the imidazo[1,2-a]pyridine, imidazo[2,1-b] thiazole, imidazotriazine, imidazo[2,1-b]-1,3,4-thiadiazole systems are described.

L13 ANSWER 36 OF 39 CAPLUS COPYRIGHT 2008 ACS on STN

Title Benzoazacycloalkanes
Author/Inventor Rossi, Alberto P. A.; Werner, Lincoln H.; Bencze, William L.; DeStevens, George
Patent Assignee/Corporate Source CIBA Ltd.
Source 55 pp.
Document Type

Patent
Language Unavailable
Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
BE 641254		19640615	BE 1964-1254	19631213

Patent Number (1)
BE 641254
Patent Publication Date (1)
19640615
Application Number (1)
BE 1964-1254
Application Date (1)
19631213
Priority Patent Number (1)
US
Priority Patent Publication Date (1)
19621214

Abstract

EtOH was slowly evaporated from a mixture of 8.9 g. 3-phenyl-2,3,4,5-tetrahydro-1-benzazepine, 2.5 g. 2-chloromethyl-2-imidazoline hydrochloride, and 30 cc. EtOH and the residue heated for 3 hrs. at 110-20° and kept overnight to give the hydrochloride of I (R1 = R2 = R3 = R4 = H, R5 = Ph, A = A' = (CH2)2, X = 1,2-phenylene) (Ia), m. 280-3°. The corresponding hydrochloride where R5 = Me was obtained by treating 1 g. Ia in 10 cc. EtOH with 1 cc. MeI, allowing the mixture to stand at ambient temperature for about 80 hrs. and evaporating in vacuo, m. 145° (decomposition). A mixture of 6.9 g. 1-cyanomethyl-3-phenyl-1,2,3,4-tetrahydroquinoline, 2.52 g. ethylenediamine, and 3 drops CS2 was heated for 4 hrs. at 130-40° and worked up to give the HCl salt of I (R1 = R2 = R3 = R4 = R5 = H, A = CH2, A' = (CH2)2, X = 1,2-phenylene) (Ia), m. 290-4° (EtOH). Similarly prepared were the following I (R2 = R3 = R4 = R5 = H, A = CH2, A' = X, and m. p. HCl salt gives): H, 2-ClC6H4, (CH2)2, (CH2)2, 1,2-phenylene (Z), 212°; H, Ph, (CH2)2, (CH2)2, 4-chloro-1,2-phenylene, 238° (decomposition); H, 4-MeC6H4, (CH2)2, (CH2)2, Z, 190-200°; H, Ph, (CH2)2, CH2CHMe, Z, 220-2°; H, Ph, (CH2)2, (CH2)2, Z, 256-9°; H, 4-ClC6H4, (CH2)2, (CH2)2, Z, 239-41°; Ph, H, CH2, (CH2)2, Z, 220°; H, 4-ClC6H4, CH2, (CH2)2, Z, 284-6°. 3-(2-Chlorophenyl)-1-cyanomethyl-2,3,4,5-tetrahydro-1-benzazepine (II) was prepared as follows: 2-chlorobenzalacetophenone (III) was prepared by treating at 15-20° an ethanolic solution of 180 g. acetophenone in aqueous NaOH with 211 g. o-chlorobenzaldehyde, the mixture shaken for 3 hrs., allowed to stand overnight, and III isolated, m. 50-2°. An ethanolic solution of 121.4 g. III containing HOAc was treated with an aqueous solution of 65 g. KCN at 55°. After cooling to 10°, 2-(o-chlorophenyl)-4-oxo-4-phenylbutyronitrile (IV) was obtained, m. 104-7° (EtOH). Methyl 2-(o-chlorophenyl)-4-oxo-4-phenylbutyrate (V) was prepared by treating an ethanolic solution of 100 g. IV with 160 g. concentrated H2SO4 and refluxing for 2.5 hrs. V was isolated on cooling, m. 105-8°. The free acid (VI), m. 137-9° (EtOH), was obtained by hydrolyzing V with NaOH. VI was then hydrogenated in the presence of a Pd catalyst at 50-60° under a pressure of about 7 atmospheric 2-(2-Chlorophenyl)-4-phenylbutyric acid (VII) was isolated, b.p. 250°. VII was cyclized by adding a benzene solution of 50 g. VII to a suspension of 44 g. PCl5 in C6H6 and heating at 100° for 30 min. The mixture was concentrated and the purified residue dissolved in C6H6 and added to a suspension of 30.4 g. anhydrous AlCl3 in C6H6 which was shaken for 2 hrs. at ambient temperature, then at 60° for 3 hrs., allowed to stand overnight, and 2-(2-chlorophenyl)-1,2,3,4-tetrahydronaphthalen-1-one (VIII), b.p. 185-90°, isolated. A compound similarly prepared was 2-(p-chlorophenyl)-1,2,3,4-tetrahydronaphthalen-1-one, m. 108-9°. A solution of 50 g. VIII in HOAc was then treated with 16.5 g. NaNH2 followed by treatment with 50 cc. concentrated H2SO4 to give 3-(2-chlorophenyl)-2,3,4,5-tetrahydro-1-benzazepin-2-one (IX), m. 240-2°. Comps. similarly prepared were 8-chloro-3-phenyl-2,3,4,5-tetrahydro-1-benzazepin-2-one, m. 224-7°; 3-(4-chlorophenyl)-2,3,4,5-tetrahydro-1-benzazepin-2-one, m. 193-5°; and 3-phenyl-2,3,4,5-tetrahydro-1-benzazepin-2-one, m. 196-8°. IX (26 g.) was then reduced with LiAlH4 in tetrahydrofuran to give 3-(2-chlorophenyl)-2,3,4,5-tetrahydro-1-benzazepine (I), m. 138-9°. Comps. similarly prepared were 8-chloro-3-phenyl-2,3,4,5-tetrahydro-1-benzazepine, m. 73-5°; 3-p-tolyl-2,3,4,5-tetrahydro-1-benzazepine, m. 76-8°; 3-(p-chlorophenyl)-2,3,4,5-tetrahydro-1-benzazepine, m. 100-2°; and 3-phenyl-2,3,4,5-tetrahydro-1-benzazepine, m. 122-4°. X (6.2 g.) was treated with an aqueous HOAc solution of 0.72 g. formaldehyde and 1.4 g. NaCN. The product was extracted with C6H6, the extract dried, and evaporated to give II. Comps. similarly prepared were 8-chloro-1-cyanomethyl-3-phenyl-2,3,4,5-tetrahydro-1-benzazepine, m. 103-4°; 3-(p-chlorophenyl)-1-cyanomethyl-2,3,4,5-tetrahydro-1-benzazepine, m. 102-4°; 3-phenyl-1-cyanomethyl-2,3,4,5-tetrahydro-1-benzazepine, b.p. 145-55°; 3-phenyl-1-cyanomethyl-1,2,3,4-tetrahydroquinoline, m. 119-20°, and 3-(p-chlorophenyl)-1-cyanomethyl-1,2,3,4-tetrahydroquinoline, m. 132-4°. 3-(p-chlorophenyl)-1,2,3,4-tetrahydroquinoline (XI) was prepared as follows: 58.5 g. N-acetylglycine was condensed with 103.6 g. p-chlorobenzaldehyde in the presence of NaOAc and Ac2 at 115-20°. After 1 hr., the mixture was cooled, filtered, and the crude 4-(p-chlorobenzylidene)-2-methyl-2-oxazolin-5-one (XII) washed with hexane and Et2O, m. 142-4°. XII was then refluxed for 4 hrs. in aqueous Me2CO to give α -acetylamino-p-chlorocinnamic acid (XIII), m. 118-20°. XIII (57.6 g.) was refluxed with N HCl for 3 hrs. to give 2-(p-chlorophenyl)pyruvic acid (XIV), m. 178-80°. Then, 60.1 g. XIV was condensed with 44.5 g. isatin in concentrated HCl at 51° to give 3-(p-chlorophenyl)quinoline-2,4-dicarboxylic acid (XV), m. 172-5° (decomposition). XV (63.7 g.) was decarboxylated by treating at 280° with 21 g. Cu powder suspended in mineral oil for 20 min. to give 3-(p-chlorophenyl)quinoline (XVI), m. 133-4° (EtOAc). XVI (15.1 g.) was hydrogenated over Raney Ni at 100° and 21.5 atmospheric to give XI, m. 106-8° (iso-PrOH).

L13 ANSWER 37 OF 39 CAPLUS COPYRIGHT 2008 ACS on STN
Title

3-Alkylthio-2-phenyl-8-(or 6)-(1-methyl-2-pyrrolyl)pyrimidazole
Author/Inventor

Godovikova, S. N.

Patent Assignee/Corporate Source

Zelinskii, N.D., Institute of Organic Chemistry, Academy of Sciences, U.S.S.R. SOURCE From: Znakov 1965(9), 25.

Byul. Izobret. i Tovarnykh

Document Type

Patent
Language Unavailable

Patent Information				
PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
SU 170513		19650423	SU	19640108

Patent Number (1)
SU 170513
Patent Publication Date (1)
19650423
Application Number (1)
SU
Application Date (1)
19640108
Priority Patent Number (1)
SU
Priority Patent Publication Date (1)
19640108

Abstract
The title compds. are prepared from the action of Br₂ in CHCl₃ on 2-phenyl-8-(or-6)-(1-methyl-2-pyrrolidyl)pyrimidazole followed by treatment of the mixture with BuLi and dialkyl disulfide.

L13 ANSWER 38 OF 39 CAPLUS COPYRIGHT 2008 ACS on STN

Title Pyridazine derivatives. VI. Synthesis of derivatives of imidazo[1,2-b]pyridazines
Author/Inventor Yoneda, Fumio; Otaka, Takayuki; Nitta, Yoshihiro
Patent Assignee/Corporate Source Chugai Pharm. A.-G., Tokyo

Source Chemical & Pharmaceutical Bulletin (1964), 12(11), 1351-6 CODEN: CPBTAL; ISSN: 0009-2363

Document Type Journal
Language German

Abstract
cf. CA 60, 10677e. The title compds. (I) were prepared for pharmacodynamic investigation, 3-Amino-6-chloropyridazine (2 g.) and 2.8 g. BzCH₂Br (II) in 100 ml. EtOH refluxed and cooled gave 3 g. I (R = Cl, R' = H) (III), m. 190° (EtOH). III (0.7 g.) in 70 ml. EtOH hydrogenated over 0.2 g. 10% Pd-C until the calculated amount H was absorbed, the solution filtered and evaporated, and the residue neutralized with aqueous NH₃ gave 0.4 g. I (R = R' = H) (IV), m. 124° (iso-Pr₂O). The following I were prepared as III (R, R' and m.p. given): Cl, OMe (V), 214°; Cl, Cl (VI), 219°; Cl, Br (VII), 222°. V, VI, and VII hydrogenated as III gave the following resp. I (same data): H, OMe, 158°, H, Cl, 183-4°; H, Br, 194-5°. 3-Aminopyridazine (1.5 g.) and 3.0 g. II in 50 ml. EtOH refluxed 2 hrs. on a water bath and evaporated gave 4 g. 1-phenacyl-6-aminopyridazinium bromide (VIII), decomposed 219° (EtOH-Et₂O). VIII (2 g.) in 50 ml. H₂O refluxed 4 hrs. and cooled gave 1.5 g. IV. HBr, decomposed 266-7° (EtOH-iso-Pr₂O), which neutralized with aqueous NH₃ gave IV, identical (mixed m.p.) with IV prepared above. III (2.1 g.) heated 5 hrs. at 120-30° with 50 ml. absolute MeOH containing 0.5 g. Na in a pressure vessel, the solution filtered and evaporated, and the residue treated with H₂O gave 2.5 g. I (R = MeO, R' = H) (VIIIa), m. 137.5° (EtOH). III (1.2 g.) and 25 ml. Me₂NCH₂CH₂OH containing 0.13 g. Na heated 3 hrs. at 130-40° and worked up as above gave 1 g. I (R = Me₂NCH₂CH₂O, R' = H), m. 107° (iso-Pr₂O). The following I were similarly prepared (same data): MeO, MeO, 182°; MeO, Cl, 179°; MeO, Br, 178°; EtO, H, 132°; EtO, MeO, 131°; EtO, Cl, 162°; EtO, Br, 171°; PrO, H, 107°; PrO, MeO, 98°; PrO, Cl, 138-9°; PrO, Br, 168°; iso-PrO, H, 98°; iso-PrO, MeO, 110°; iso-PrO, Cl, 138-9°; iso-PrO, Br, 148-9°; Me₂NCH₂CH₂O, MeO, 103-4°; Me₂NCH₂CH₂O, Cl, 158°; Me₂NCH₂CH₂O, Br, 172°; Et₂NCH₂CH₂O, H, 102°; Et₂NCH₂CH₂O, MeO, 75°; Et₂NCH₂CH₂O, Cl, 129°; Et₂NCH₂CH₂O, Br, 132.5°. III (1.5 g.) in 50 ml. EtOH containing excess Me₂NH heated 5 hrs. at 170-80° in a pressure vessel and evaporated gave 1.5 g. I (R = Me₂N, R' = H), m. 195° (EtOH). The following I were similarly prepared (same data): Me₂N, MeO, 204°; Me₂N, Cl, 203°; Me₂N, Br, 197°; morpholino, H, 183-4°; morpholino, MeO, 198-9°; morpholino, Cl, 227°; morpholino, Br, 234°; piperidino, H, 169°; piperidino, MeO, 153°; piperidino, Cl, 222°; piperidino, Br, 221°; pyrrolidino, H, 187-8°; pyrrolidino, MeO, 190°; pyrrolidino, Cl, 210°; pyrrolidino, Br, 211°. III (6 g.) in 100 ml. 90% EtOH containing 6 g. KOH heated 4 hrs. at 160-70° in a pressure vessel, cooled, and evaporated, the residue treated with H₂O, and the solution filtered and neutralized with aqueous HCl gave 5.5 g. I (R = HO, R' = H) (IX), m. 300° (EtOH). Similarly were prepared the following I (same data): HO, MeO, 287°; HO, Cl, 300°; HO, Br, 300°. IX (0.65 g.) in 20 ml. MeOH containing 0.21 g. KOH refluxed 2 hrs. with 1.03 g. MeI, the solution concentrated, and the residue treated with H₂O gave 0.7 g. X (R = H), m. 120° (EtOH-H₂O). Similarly were prepared the following X (R and m.p. given): MeO, 140-1°; Cl, 246-7°; Br, 261-2°. The uv spectra of IV, VIIIa, IX, and X (R = H) were recorded, as well as that of 2-phenylimidazo[1,2-a]pyridine (for comparison with IV).

L13 ANSWER 39 OF 39 CAPLUS COPYRIGHT 2008 ACS on STN

Title 8-(and 6)-(1-methyl-2-pyrrolidyl)-2-phenylimidazo[1,2-a]pyridine
Author/Inventor Gol'dfarb, Ya. L.; Andrichuk, M. V.
Document Type Patent
Language

Unavailable

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
RU 51041		19370531	RU	

Patent Number (1)

RU 51041

Patent Publication Date (1)

19370531

Application Number (1)

RU

Abstract

α - or α' -Aminonicotine is heated in the presence of a solvent with α -bromoacetophenone. Cf. C. A. 32, 176.6.

->

->



ring nodes :

1 2 3 4 5 6 7 8 9

ring bonds :

1-2 1-5 1-9 2-3 3-4 4-5 5-6 6-7 7-8 8-9

exact/norm bonds :

1-2 1-5 1-9 2-3 3-4 4-5 5-6 6-7 7-8 8-9

G1:C,N

Match level :

1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:Atom 8:Atom 9:Atom

L14 STRUCTURE UPLOADED

=> g l14 sss sam

SAMPLE SEARCH INITIATED 11:21:15 FILE 'REGISTRY'

SAMPLE SCREEN SEARCH COMPLETED - 11451 TO ITERATE

17.5% PROCESSED 2000 ITERATIONS

50 ANSWERS

INCOMPLETE SEARCH (SYSTEM LIMIT EXCEEDED)

SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS: ONLINE **COMPLETE**

BATCH **COMPLETE**

PROJECTED ITERATIONS: 222608 TO 235432

PROJECTED ANSWERS: 78820 TO 86532

L15 50 SEA SSS SAM L14

=> s l14 sss full
FULL SEARCH INITIATED 11:21:23 FILE 'REGISTRY'
FULL SCREEN SEARCH COMPLETED - 228722 TO ITERATE

100.0% PROCESSED 228722 ITERATIONS 80930 ANSWERS
SEARCH TIME: 00.00.01

L16 80930 SEA SSS FUL L14

=> s l16
L17 6936 L16

=> s l17 and electroluminescent
68951 ELECTROLUMINESCENT
6 ELECTROLUMINESCENTS
68953 ELECTROLUMINESCENT
(ELECTROLUMINESCENT OR ELECTROLUMINESCENTS)

L18 42 L17 AND ELECTROLUMINESCENT

=> s l18 and py<-2003
23975208 PY<-2003

L19 21 L18 AND PY<-2003

=> d l19 1-21 ibib abs

L19 ANSWER 1 OF 21 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Carbazole derivative for organic electroluminescent devices and organic electroluminescent devices

Author/Inventor

Iwakuma, Toshihiro; Yamamoto, Hiroshi; Hironaka, Yoshio; Ikeda, Hidetsugu; Hosokawa, Chishio; Tomita, Seiji; Arakane, Takashi

Patent Assignee/Corporate Source

Idemitsu Kosan Co., Ltd., Japan

Source

PCT Int. Appl., 68 pp. CODEN: PIXXD2

Document Type

Patent

Language

Japanese

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003080760	A1	20031002	WO 2003-JP3329	20030319

Patent Number (1)

WO 2003080760

Kind Code (1)

A1

Patent Publication Date (1)

20031002

Application Number (1)

WO 2003-JP3329

Application Date (1)

20030319

Priority Patent Number (1)

JP 2002-81234

Priority Kind Code (1)

A

Priority Patent Publication Date (1)

20020322

Abstract

The invention refers to a material for blue electroluminescent devices having the structure (Cz)_nA or Cz(A)_n [Cz = (un)substituted arylcarbazolyl or carbazoylalkylene; A = MpLqM'r; M,M' = (un)substituted C2-40 heteroarom. rings; L = single bond, (un)substituted C6-30 aryl or arylene, C5-30 cycloalkylene, photorefractive C2-30 heteroarom.; p,r = 0 - 2; q = 1 - 2; p + r > 1].

L19 ANSWER 2 OF 21 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Blue-emitting organic electroluminescent devices of high color purity and materials therefor

Author/Inventor

Iwakuma, Toshihiro; Yamamoto, Hiroshi; Hironaka, Yoshio; Ikeda, Shuji; Hosokawa, Chishio

Patent Assignee/Corporate Source

Idemitsu Kosan Co., Ltd., Japan

Source

Jpn. Kokai Tokkyo Koho, 15 pp. CODEN: JKXXAF

Document Type
Patent
Language

Japanese

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2003277743	A	20031002	JP 2002-80817	20020322

Patent Number (1)

JP 2003277743

Kind Code (1)

A

Patent Publication Date (1)

20031002

Application Number (1)

JP 2002-80817

Application Date (1)

20020322

Priority Patent Number (1)

JP 2002-80817

Priority Patent Publication Date (1)

20020322

Abstract

Comps. (Ind)ⁿA or IndAm [Ind = arylindole or indolealkylene; A = MpLqM^r (M, M^r = C2-40 azacyclic aromatic ring; L = single bond, C6-30 aryene, C5-30 cycloalkylene, C2-30 aromatic heterocycle; p = 0-2; q = 1-2; r = 0-2; p + r ≥ 1)] are claimed. Blue-emitting electroluminescent devices including (emission) layers containing the comps. as host materials are also claimed.

L19 ANSWER 3 OF 21 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Materials for organic blue electroluminescent device for organic electroluminescent display

Author/Inventor

Iwakuma, Toshihiro; Hosokawa, Chishio; Yamamoto, Hiroshi

Patent Assignee/Corporate Source

Idemitsu Kosan Co., Ltd., Japan

Source

Jpn. Kokai Tokkyo Koho, 10 pp. CODEN: JKXXAF

Document Type
Patent
Language

Japanese

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2003268362	A	20030925	JP 2002-71397	20020315

Patent Number (1)

JP 2003268362

Kind Code (1)

A

Patent Publication Date (1)

20030925

Application Number (1)

JP 2002-71397

Application Date (1)

20020315

Priority Patent Number (1)

JP 2002-71397

Priority Patent Publication Date (1)

20020315

Abstract

The title material contains compound (Ind)ⁿmM (Ind = indole; M = C2-40 heterocyclic ring containing N; n, m = 1-3 integer). The device provides blue with good color purity.

L19 ANSWER 4 OF 21 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Double-spiro organic compounds and electroluminescent devices

Author/Inventor

Kim, Kong-Kyeum; Son, Se-Hwan; Yoon, Seok-Hee; Bae, Jae-Soon; Lee, Youn-Gu; Im, Sung-Gap; Kim, Ji-Eun; Lee, Jae-Chol

Patent Assignee/Corporate Source

LG Chem, Ltd., S. Korea

Source

PCT Int. Appl., 117 pp. CODEN: PIXXD2

Document Type

Patent

Language

English

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002088274	A1	20021107	WO 2002-KR458	20020318

Patent Number (1)

WO 2002088274

Kind Code (1)

A1

Patent Publication Date (1)

20021107

Application Number (1)

WO 2002-KR458

Application Date (1)

20020318

Priority Patent Number (1)

KR 2001-23038

Priority Kind Code (1)

A

Priority Patent Publication Date (1)

20010427

Abstract

Double-spiro organic compds. are claimed which are described by the general formula I (R1-24 = independently selected substituents not all of which are H). Light-emitting, hole-transporting, and electron-transporting materials comprising the compds. are also described. Electroluminescent materials comprising the compds. including deposited films, methods for depositing the materials, and organic electroluminescent devices employing the materials, and method for fabricating the devices, are also described.

L19 ANSWER 5 OF 21 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Red-emitting organic electroluminescent devices with high electric energy conversion efficiency and color purity

Author/Inventor

Tominaga, Takeshi; Murase, Seichiro; Kohama, Toru

Patent Assignee/Corporate Source

Toray Industries, Inc., Japan

Source

Jpn. Kokai Tokkyo Koho, 13 pp. CODEN: JKXXAF

Document Type

Patent

Language

Japanese

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2002008862	A	20020111	JP 2000-184268	20000620

Patent Number (1)

JP 2002008862

Kind Code (1)

A

Patent Publication Date (1)

20020111

Application Number (1)

JP 2000-184268

Application Date (1)

20000620

Priority Patent Number (1)

JP 2000-184268

Priority Patent Publication Date (1)

20000620

Abstract

The devices having emission peak at 580-750 nm, contain fluorescent substances having fluorescent peak at 540-750 nm and condensed heterocyclic compds. (which may be dopants), between anodes and cathodes. The compds. may have polar groups, vinyl groups, aromatic rings, and/or heterocyclic rings. The devices are useful for matrix-type displays (e.g., computers, televisions) and segment-type displays (e.g., clocks, thermometers).

L19 ANSWER 6 OF 21 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Organic electroluminescent devices using condensed heterocyclic rings

Author/Inventor

Taguchi, Toshiaki; Mishima, Masayuki; Ise, Toshihiro; Okada, Hisashi
Patent Assignee/Corporate Source
Fuji Photo Film Co., Ltd., Japan

Source
Jpn. Kokai Tokkyo Koho, 17 pp. CODEN: JKXXAF

Document Type
Patent

Language
Japanese

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2001357977	A	20011226	JP 2000-175981	20000612

Patent Number (1)

JP 2001357977

Kind Code (1)

A

Patent Publication Date (1)

20011226

Application Number (1)

JP 2000-175981

Application Date (1)

20000612

Priority Patent Number (1)

JP 2000-175981

Priority Patent Publication Date (1)

20000612

Abstract

The invention relates to an organic electroluminescent device comprising a pair of electrodes sandwiching ≥ 1 layer(s) containing ≥ 1 condensed heterocyclic compds. represented by I (X = C, N; Z1 and Z2 may form a N-containing heterocyclyl) and ortho metal (Ir) complexes. The device shows high luminance, luminescent efficiency, and superior in durability.

L19 ANSWER 7 OF 21 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Electroluminescent device using condensed rings

Author/Inventor

Kohama, Toru; Tominaga, Takeshi; Murase, Seiichiro

Patent Assignee/Corporate Source

Toray Industries, Inc., Japan

Source

Jpn. Kokai Tokkyo Koho, 12 pp. CODEN: JKXXAF

Document Type

Patent

Language

Japanese

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2001338764	A	20011207	JP 2000-159883	20000530

Patent Number (1)

JP 2001338764

Kind Code (1)

A

Patent Publication Date (1)

20011207

Application Number (1)

JP 2000-159883

Application Date (1)

20000530

Priority Patent Number (1)

JP 2000-159883

Priority Patent Publication Date (1)

20000530

Abstract

The invention relates to a red-emitting electroluminescent device having the emission peak in 580 - 720 nm between the anode and the cathode, wherein the electroluminescent layer comprises the fluorescent substance having the emission peak in 540 - 720 nm as a host material, and the condensed rings I, II, III [R1-46 = the same or different groups selected from H, alkyl, alkoxy, halo, aryl, alkaryl, alkenyl, aryl ether, heterocyclyl, cyano, aldehyde, CO, ester, carbamoyl, amino and fused rings or aliphatic rings formed with adjacent substituents]. The red luminous component offers superior in color purity.

L19 ANSWER 8 OF 21 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Electroluminescent device using coumarin derivatives

Author/Inventor
Kohama, Toru; Tominaga, Takeshi; Murase, Seichiro
Patent Assignee/Corporate Source
Toray Industries, Inc., Japan
Source
Jpn. Kokai Tokkyo Koho, 10 pp. CODEN: JKXXAF
Document Type
Patent
Language
Japanese

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2001338763	A	20011207	JP 2000-159882	20000530

Patent Number (1)
JP 2001338763
Kind Code (1)
A
Patent Publication Date (1)
20011207
Application Number (1)
JP 2000-159882
Application Date (1)
20000530
Priority Patent Number (1)
JP 2000-159882
Priority Patent Publication Date (1)
20000530

Abstract

The invention relates to a red-emitting electroluminescent device having the emission peak in 580 - 720 nm between the anode and the cathode, wherein the electroluminescent layer comprises the fluorescent substance having the emission peak in 540 - 720 nm as a host material, and the coumarins I [R1-6 = the same or different groups selected from H, alkyl, alkoxy, halo, aryl, aralkyl, alkenyl, aryl ether, heterocycl, cyano, aldehyde, CO, ester, carbamoyl, amino and fused rings or aliphatic rings formed with adjacent substituents]. The red luminous component offers superior in color purity.

L19 ANSWER 9 OF 21 CAPLUS COPYRIGHT 2008 ACS ON STN

Title

Organic electroluminescent device

Author/Inventor
Kohama, Toru; Tominaga, Takeshi; Murase, Seichiro
Patent Assignee/Corporate Source
Toray Industries, Inc., Japan
Source
Jpn. Kokai Tokkyo Koho, 11 pp. CODEN: JKXXAF
Document Type
Patent
Language
Japanese

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2001313175	A	20011109	JP 2000-129396	20000428

Patent Number (1)
JP 2001313175
Kind Code (1)
A
Patent Publication Date (1)
20011109
Application Number (1)
JP 2000-129396
Application Date (1)
20000428
Priority Patent Number (1)
JP 2000-129396
Priority Patent Publication Date (1)
20000428

Abstract

The invention relates to a red-emitting organic electroluminescent device having the emission peak in 580-720 nm, suited for use in making segment- and matrix-type displays, a backlight, an illumination apparatus, etc., wherein the electroluminescent layer comprises the fluorescent substance having the emission peak in 540-720 nm, as a host material, and the polycyclic ketone represented by I and II [R1-17 = H, alkyl, alkoxy, halo, etc.], as a dopant.

L19 ANSWER 10 OF 21 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Organometallic electroluminescent devices

Author/Inventor

Kitasawa, Daisuke; Tominaga, Takeshi; Takano, Akiko

Patent Assignee/Corporate Source

Toray Industries, Inc., Japan

Source

Jpn. Kokai Tokkyo Koho, 9 pp. CODEN: JKXXAF

Document Type

Patent

Language

Japanese

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2001057292	A	20010227	JP 1999-233791	19990820

Patent Number (1)

JP 2001057292

Kind Code (1)

A

Patent Publication Date (1)

20010227

Application Number (1)

JP 1999-233791

Application Date (1)

19990820

Priority Patent Number (1)

JP 1999-233791

Priority Patent Publication Date (1)

19990820

Abstract

The devices comprise a phosphor containing a metal complex having ligands comprising an imidazopyridine derivative I (R1-6 = H, alkyl, cycloalkyl, aralkyl, alkenyl, cycloalkenyl, OH, mercapto, alkoxy, alkylthio, arylether, arylthioether, aryl, heterocyclic, halo, haloalkane, haloalkene, CN, aldehyde, carbonyl, carboxyl, ester, carbo moi, amino, nitro, silyl, siloxanyl).

L19 ANSWER 11 OF 21 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Organic electroluminescent device

Author/Inventor

Nakatsuka, Masakatsu; Shimamura, Takehiko

Patent Assignee/Corporate Source

Mitsui Chemicals Inc., Japan

Source

Jpn. Kokai Tokkyo Koho, 43 pp. CODEN: JKXXAF

Document Type

Patent

Language

Japanese

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2001035664	A	20010209	JP 1999-206147	19990721

Patent Number (1)

JP 2001035664

Kind Code (1)

A

Patent Publication Date (1)

20010209

Application Number (1)

JP 1999-206147

Application Date (1)

19990721

Priority Patent Number (1)

JP 1999-206147

Priority Patent Publication Date (1)

19990721

Abstract

The invention relates to an organic electroluminescent device comprising an azaindolizine-containing layer placed between a pair of electrodes.

L19 ANSWER 12 OF 21 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Organic electroluminescent devices

Author/Inventor

Tominaga, Takeshi; Kohama, Toru; Takano, Akiko

Patent Assignee/Corporate Source

Toray Industries, Inc., Japan

Source

Jpn. Kokai Tokkyo Koho, 20 pp. CODEN: JKXXAF

Document Type

Patent

Language

Japanese

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2001006877	A	20010112	JP 1999-174058	19990621

Patent Number (1)

JP 2001006877

Kind Code (1)

A

Patent Publication Date (1)

20010112

Application Number (1)

JP 1999-174058

Application Date (1)

19990621

Priority Patent Number (1)

JP 1999-174058

Priority Patent Publication Date (1)

19990621

Abstract

The devices comprise an imidazopyridine phosphor I (R1-6 = H, alkyl, cycloalkyl, aralkyl, alkenyl, cycloalkenyl, alkoxy, alkylthio, aryl ether, aryl thioether, aryl, heterocyclic, halo, cyano, aldehyde, carbonyl, carboxyl, ester, carbamoyl, amino, nitro, silyl, siloxanyl).

L19 ANSWER 13 OF 21 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Electroluminescent element and devices

Author/Inventor

Kohama, Akira; Tominaga, Tsuyoshi; Kitazawa, Daisuke; Himeshima, Yoshio

Patent Assignee/Corporate Source

Toray Industries, Inc., Japan

Source

Eur. Pat. Appl., 22 pp. CODEN: EPXXDW

Document Type

Patent

Language

English

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 1000998	A1	20000517	EP 1999-308823	19991105

Patent Number (1)

EP 1000998

Kind Code (1)

A1

Patent Publication Date (1)

20000517

Application Number (1)

EP 1999-308823

Application Date (1)

19991105

Priority Patent Number (1)

JP 1998-317681

Priority Kind Code (1)

A

Priority Patent Publication Date (1)

19981109

Abstract

Electroluminescent devices are described which emit a peak wavelength at ≤ 580 nm in which the active material contains at least a fluorescent compound with a fluorescence maximum at ≥ 540 nm or above and a compound with a pyromethene structure described by the general formula I or a metal complex thereof (R1-7 = the same or different groups selected from H, alkyl, alkoxy,

halo, aryl, alkyl, alkenyl, aryl ether, heterocyclic, cyano, aldehyde, CO, ester, carbamoyl, amino and fused rings or aliphatic rings formed with adjacent substituents; and X = C or nitrogen, with the restriction that, where X = nitrogen, R7 is absent). Display, signaling and illumination devices employing the elements are described.

L19 ANSWER 14 OF 21 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Organic Pigment Nanoparticle Thin Film Devices via Lewis Acid Pigment Solubilization

Author/Inventor

Hsieh, Bing R.; Melnyk, Andrew R.

Patent Assignee/Corporate Source

Xerox Corporation, Webster, NY, 14580, USA

Source

Chemistry of Materials (1998), 10(9), 2313-2316 CODEN: CMATEX; ISSN: 0897-4756

Document Type

Journal

Language

English

Abstract

Lewis acid pigment solubilization (LAPS) deposition of thin films of pigment particles is useful for fabrication of pigment-based organic semiconductor devices like light-emitting diodes, photodiodes, solar cells and field effect transistors. The process involves the solubilization of a pigment in a Lewis acid/nitromethane solution, and casting the pigment/Lewis acid thin films on substrates with a predeposited adhesive layer. The authors applied the LAPS process for fabrication of multilayer electrophotog. organic photoconductors. A mixture containing bis(benzimidazole)perylene (BZP), AlCl₃, nitromethane, and methylene chloride was stirred under N₂ and coated on a metalized Mylar support overcoated with a thin 2-aminopropyltriethoxysilane primer layer and DuPont 46K polyester adhesive layer. The molar ratio of BZP to AlCl₃ was 1:6, solns. having 2 weight% BZP were easily filtered through a 0.45 µm filter, but solns. with higher pigment concns. could not be easily filtered. The as-cast BZP/AlCl₃ layers were washed with water, followed by wash with aqueous Na₂CO₃ (or NH₄OH) to produce charge-generation layer. The transport layer was deposited by draw bar coating of a solution containing N,N'-diphenyl-N,N'-(m-tolyl)-1,1'-biphenyl-4,4'-diamine, Macrolon and methylene chloride. The pigment can also be dispersed in an electrophotocative polymer binders. Xerog. properties of the photoconductors prepared by LAPS method had good photosensitivity, high cyclic stability, low dark decay and residual charges.

L19 ANSWER 15 OF 21 CAPLUS COPYRIGHT 2008 ACS on STN

Title

An organic light emitting device containing a protection layer

Author/Inventor

Thompson, Mark E.; Forrest, Stephen R.; Burrows, Paul; Garbuzov, Dmitri Z.; Shen, Zilan; Cronin, Jon A.; et al.

Patent Assignee/Corporate Source

The Trustees of Princeton University, USA; The University of Southern California

Source

PCT Int. Appl., 107 pp. CODEN: PIXXD2

Document Type

Patent

Language

English

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9828767	A1	19980702	WO 1997-US23952	19971223

Patent Number (1)

WO 9828767

Kind Code (1)

A1

Patent Publication Date (1)

19980702

Application Number (1)

WO 1997-US23952

Application Date (1)

19971223

Priority Patent Number (1)

US 1996-771815

Priority Kind Code (1)

A

Priority Patent Publication Date (1)

19961223

Abstract

Organic light-emitting devices comprising a heterostructure for producing electroluminescence are described in which the heterostructure includes a protection layer between a hole-transporting layer and an (e.g., indium tin oxide) anode layer; devices incorporating down-conversion layers are also described. Methods of fabricating the devices are provided, and the use of the devices in displays is described.

L19 ANSWER 16 OF 21 CAPLUS COPYRIGHT 2008 ACS on STN

Title Exchanging light and charges in electronic polymers
 Author/Inventor Jenekhe, Samson A.; Zhang, Xuejun
 Patent Assignee/Corporate Source Dep. Chemical Eng., Univ. Rochester, Rochester, NY, 14627, USA
 Source Annual Technical Conference - Society of Plastics Engineers (1996), 54th(Vol. 2), 1323-1326 CODEN: ACPED4; ISSN: 0272-5223
 Document Type Journal
 Language English

Abstract Optoelectronic devices such as xerog. photoreceptors (PRs) and light emitting diodes (LEDs) represent two examples of promising applications of semiconducting polymers. In the former example, efficient generation of charge carriers from absorbed light is the central focus. In the latter examples, efficient generation of light from injected charges is the goal. To explore the nature of these light ↔ charges exchange processes in electronic polymers we have fabricated and evaluated bilayer photoreceptors and bilayer LEDs from conjugated polymers. The efficiencies and other performance characteristics of these devices are shown to depend not only on mol. structure and intrinsic properties of the materials but also on the "polymer device engineering" factors such as the polymer/polymer interfaces.

L19 ANSWER 17 OF 21 CAPLUS COPYRIGHT 2008 ACS on STN

Title Organic electroluminescent device
 Author/Inventor Kang, Wen-Bing; Yu, Nu; Tokida, Akihiko; Potrawa, Thomas; Winterfeldt, Andreas
 Patent Assignee/Corporate Source Hoechst A.-G., Germany
 Source PCT Int. Appl., 19 pp. CODEN: PIXXD2
 Document Type Patent
 Language English

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9605267	A1	19960222	WO 1995-EP3128	19950807

Patent Number (1)
 WO 9605267
 Kind Code (1)
 A1
 Patent Publication Date (1)
 19960222
 Application Number (1)
 WO 1995-EP3128
 Application Date (1)
 19950807
 Priority Patent Number (1)
 JP 1994-185820
 Priority Kind Code (1)
 A
 Priority Patent Publication Date (1)
 19940808

Abstract Organic electroluminescent devices comprising an anode, and organic hole transport layer, an organic emission layer, an optional electron transport layer, and a cathode layered in this order on a substrate, are described in which the organic emission layer comprises a naphthalimide derivative described by the formula I (R1 = a hydrogen atom, a straight or branched chain C1-16 alkyl group, and aryl group which may have ≥1 substituents or an aralkyl group which may have ≥1 substituents; R2, R3, R4 = independently selected C1-16 alkyl, C1-16 alkoxy, C1-16 ester, optionally substituted C4-10 aryl, or optionally substituted C4-10 aryloxy groups; -CN, -CF3, -F, or -NR5R6; R5, R6 = independently selected C1-16 alkyl groups or optionally substituted C4-10 aryl groups; and m, n, p = 0, 1, 2, or 3).

L19 ANSWER 18 OF 21 CAPLUS COPYRIGHT 2008 ACS on STN

Title Conjugated polymer exciplexes and applications thereof
 Author/Inventor Jenekhe, Samson A.; Osaheni, John A.
 Patent Assignee/Corporate Source Research Corp. Technologies, Inc., USA
 Source PCT Int. Appl., 188 pp. CODEN: PIXXD2

Document Type

Patent

Language

English

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9512628	A1	19950511	WO 1994-US12322	19941028

Patent Number (1)

WO 9512628

Kind Code (1)

A1

Patent Publication Date (1)

19950511

Application Number (1)

WO 1994-US12322

Application Date (1)

19941028

Priority Patent Number (1)

US 1993-146266

Priority Kind Code (1)

A

Priority Patent Publication Date (1)

19931102

Abstract

Exciplexes with good luminescence, photogeneration of charge carriers, and quantum efficiency are formed from a π -conjugated polymers such as poly(p-phenylenebenzobisoxazole) (I) and an electron donor or acceptor component. Also claimed are assemblies comprising these exciplexes, their use in optoelec. devices and a method of enhancing optoelec. properties of π -conjugated polymers by forming these exciplexes. A typical H-tris(p-tolyl)amine (II) exciplex was prepared by spin-coating a MeNO₂ solution of I and AlCl₃ onto glass and fused silica substrates and overcoating with a CH₂Cl₂ solution containing a 40:60 II-bisphenol A polycarbonate mixture

L19 ANSWER 19 OF 21 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Organic photoreceptors and their film light-emitting devices

Author/Inventor

Maruyama, Shigeru; Kosho, Noboru; Origasa, Hitoshi; Kawate, Kenji

Patent Assignee/Corporate Source

Fuji Electric Co Ltd, Japan

Source

Jpn. Kokai Tokkyo Koho, 27 pp. CODEN: JKXXAF

Document Type

Patent

Language

Japanese

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 05134430	A	19930528	JP 1991-297513	19911114

Patent Number (1)

JP 05134430

Kind Code (1)

A

Patent Publication Date (1)

19930528

Application Number (1)

JP 1991-297513

Application Date (1)

19911114

Priority Patent Number (1)

JP 1991-297513

Priority Patent Publication Date (1)

19911114

Abstract

The title organic optical device utilize a charge/light-emission layer and a charge injection transporting layer; the formation layer effects corrosion between charge (pos. hole and electron pair) and light with light emission (from the light-generating material) resulting from the injection of charge from the charge-transporting layer and charge generation resulting from exposure to light, and the latter layer comprising a charge-transporting material and a poly(phenylene oxide) binder simultaneous to effecting charge transfer with the charge/light-conversion layer effects the flow of charge under a potential gradient. Polycarbonate may also be used as binder, and the charge-transporting material may be a diamine, hydrazone, or a stilbene derivative

L19 ANSWER 20 OF 21 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Organic electroluminescent devices

Author/Inventor

Utsuki, Koji; Nagahata, Emi

Patent Assignee/Corporate Source

Nippon Electric Co, Japan

Source

Jpn. Kokai Tokkyo Koho, 5 pp. CODEN: JKXXAF

Document Type

Patent

Language

Japanese

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 05041282	A	19930219	JP 1991-196341	19910806

Patent Number (1)

JP 05041282

Kind Code (1)

A

Patent Publication Date (1)

19930219

Application Number (1)

JP 1991-196341

Application Date (1)

19910806

Priority Patent Number (1)

JP 1991-196341

Priority Patent Publication Date (1)

19910806

Abstract

The device comprises : an electron-injecting electrode; an optional organic electron-transport layer (oxidation potential $X = X1$); an organic phosphor ($X = X2$); an organic hole-transport layer ($X = X3$); and a transparent electrode-formed substrate, wherein $X1 > X2$, $X1 > X3$, and $X3 > X2$. An alternative device comprises an organic phosphor ($X = X4$, reduction potential $Y = Y4$) which is dispersed in an electron-transport layer ($X = X5$, $Y = Y5$), wherein $X4 > X5$ and $Y4 > Y5$. The devices having a low threshold exhibit a high conversion efficiency.

L19 ANSWER 21 OF 21 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Organic thin-film electroluminescent elements

Author/Inventor

Utsuki, Koji; Takahashi, Hiroshi

Patent Assignee/Corporate Source

NEC Corp., Japan; Nippon Soda Co., Ltd.

Source

Jpn. Kokai Tokkyo Koho, 6 pp. CODEN: JKXXAF

Document Type

Patent

Language

Japanese

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 04110390	A	19920410	JP 1990-232100	19900831

Patent Number (1)

JP 04110390

Kind Code (1)

A

Patent Publication Date (1)

19920410

Application Number (1)

JP 1990-232100

Application Date (1)

19900831

Priority Patent Number (1)

JP 1990-232100

Priority Patent Publication Date (1)

19900831

Abstract

The element comprises: (1) a transparent substrate, (2) a transparent electrode, (3) an optional hole-transport, (4) a phosphor, (5) an optional electron-transport, and (6) a (transparent) electrode layer, wherein (2)-(5) employ organic materials; and (4) consists of a pyrimidazoquinoxaline derivative I(R1-8=H, halo, OH, NO₂, (un)substituted-alkyl, -aryl, -pyridyl, -benzoyl). The element provides a

luminous yellow-emitting defect-free light source for large-area flat displays.

=>

Connecting via Winsock to STN

Welcome to STN International! Enter x:X


LOGINID:SSPTAMEN1774

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

***** Welcome to STN International *****

=>



chain nodes :

27

ring nodes :

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26


```

chain bonds :
7-21 10-15 24-27

ring bonds :
1-2 1-6 2-3 3-4 4-5 5-6 5-7 6-10 7-8 8-9 8-11 9-10 9-14 11-12 12-13 13-14 15-16 15-20 16-17 17-18 18-19
19-20 21-22 21-26 22-23 23-24 24-25 25-26

exact/norm bonds :
21-22 21-26 22-23 23-24 24-25 24-27 25-26

exact bonds :
7-21 10-15

normalized bonds :
1-2 1-6 2-3 3-4 4-5 5-6 5-7 6-10 7-8 8-9 8-11 9-10 9-14 11-12 12-13 13-14 15-16 15-20 16-17 17-18 18-19
19-20

Match level :
1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:Atom 8:Atom 9:Atom 10:Atom 11:Atom 12:Atom 13:Atom 14:Atom
15:Atom 16:Atom 17:Atom 18:Atom 19:Atom 20:Atom 21:Atom 22:Atom 23:Atom 24:Atom 25:Atom 26:Atom
27:Atom

```

```

L1      STRUCTURE UPLOADED

=> s l1 sss sam
SAMPLE SEARCH INITIATED 16:44:51 FILE 'REGISTRY'
SAMPLE SCREEN SEARCH COMPLETED -      3250 TO ITERATE

61.5% PROCESSED      2000 ITERATIONS      29 ANSWERS
INCOMPLETE SEARCH (SYSTEM LIMIT EXCEEDED)
SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS:  ONLINE      **COMPLETE**
                        BATCH      **COMPLETE**
PROJECTED ITERATIONS:    61581 TO    68419
PROJECTED ANSWERS:       531 TO     1353

L2      29 SEA SSS SAM L1

=> s l1 sss full
FULL SEARCH INITIATED 16:44:56 FILE 'REGISTRY'
FULL SCREEN SEARCH COMPLETED -    66509 TO ITERATE

100.0% PROCESSED    66509 ITERATIONS      762 ANSWERS
SEARCH TIME: 00.00.01

L3      762 SEA SSS FUL L1

=> s l3
L4      262 L3

=> s l4 and (electroluminescent or electroluminescence)
68951 ELECTROLUMINESCENT
6      ELECTROLUMINESCENTS
68953 ELECTROLUMINESCENT
      (ELECTROLUMINESCENT OR ELECTROLUMINESCENTS)
22377 ELECTROLUMINESCENCE
25     ELECTROLUMINESCENCES
22381 ELECTROLUMINESCENCE
      (ELECTROLUMINESCENCE OR ELECTROLUMINESCENCES)
L5      186 L4 AND (ELECTROLUMINESCENT OR ELECTROLUMINESCENCE)

```

=> s 15 and py<-2003
23975208 PY<-2003
L6 37 L5 AND PY<-2003
-> d 16 1-37 ibib abs

L6 ANSWER 1 OF 37 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Organic light-emitting diode devices with improved operational stability

Author/Inventor

Jarikov, Viktor V.

Patent Assignee/Corporate Source

Eastman Kodak Company, USA

Source

U.S. Pat. Appl. Publ., 108 pp., Cont.-in-part of U.S. Ser. No. 131,801, abandoned. CODEN: USXXCO

Document Type

Patent

Language

English

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2004076853	A1	20040422	US 2003-634324	20030805

Patent Number (1)

US 2004076853

Kind Code (1)

A1

Patent Publication Date (1)

20040422

Application Number (1)

US 2003-634324

Application Date (1)

20030805

Priority Patent Number (1)

US 2002-131801

Priority Kind Code (1)

B2

Priority Patent Publication Date (1)

20020424

Abstract

Organic light-emitting devices which comprise a substrate; an anode and a cathode disposed over the substrate; a luminescent layer disposed between the anode and the cathode are described in which the luminescent layer includes a host and ≥ 1 dopant; the host including a solid organic material comprising a mixture of ≥ 2 components including a first component that is an organic compound capable of transporting either electrons and/or holes and of forming both monomer state and an aggregate state and a second component of that is an organic compound that upon mixing with the first host component is capable of forming a continuous and substantially pin-hole-free layer, while the dopant of is selected to produce light from the light-emitting device. The first component is capable of forming an aggregate state either in the ground electronic state or in an excited electronic state that results in a different absorption or emission spectrum or both relative to the absorption or emission spectrum or both of the monomer state, resp., or of forming an aggregate state whose presence results in a quantum yield of luminescence of the monomer state being different relative to the quantum yield of luminescence of the monomer state in the absence of the aggregate state. The aggregate state may be crystalline

L6 ANSWER 2 OF 37 CAPLUS COPYRIGHT 2008 ACS on STN

Title

New organic compounds for electroluminescence and organic electroluminescent devices using the same

Author/Inventor

Kim, Ji-Eun; Son, Se-Hwan; Bae, Jae-Soon; Lee, Youn-Gu; Kim, Kong-Kyeum; Lee, Jae-Chol; Jang, Jun-Gi; Im, Sung-Gap

Patent Assignee/Corporate Source

LG Chem, Ltd., S. Korea

Source

PCT Int. Appl., 145 pp. CODEN: PIXXD2

Document Type

Patent

Language

English

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003095445	A1	20031120	WO 2003-KR899	20030506

Patent Number (1)

WO 2003095445

Kind Code (1)

A1
 Patent Publication Date (1)
 20031120
 Application Number (1)
 WO 2003-KR899
 Application Date (1)
 20030506
 Priority Patent Number (1)
 KR 2002-25084
 Priority Kind Code (1)
 A
 Priority Patent Publication Date (1)
 20020507

Abstract

Disclosed is a novel group of compds. having a general structure of anthracene body substituted with at least one thiophenyl group, which can be further substituted with various substituent groups. These new compds. are generally compatible with organic electroluminescence. Also disclosed are organic electroluminescent devices and method of making the same. The organic electroluminescent devices include at least one of the compds. in various layers thereof. Organic electroluminescent devices employing the new compds. in their light-emitting layers show outstanding stability.

L6 ANSWER 3 OF 37 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Electroluminescent devices

Author/Inventor

Xie, Shuang

Patent Assignee/Corporate Source

Can.

Source

U.S. Pat. Appl. Publ., 32 pp. CODEN: USXXCO

Document Type

Patent

Language

English

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2003215667	A1	20031120	US 2001-985204	20011102

Patent Number (1)

US 2003215667

Kind Code (1)

A1

Patent Publication Date (1)

20031120

Application Number (1)

US 2001-985204

Application Date (1)

20011102

Priority Patent Number (1)

US 2001-985204

Priority Patent Publication Date (1)

20011102

Abstract

Organic electroluminescent devices are described which are provided with active layers comprising a host based on ≥ 1 anthracene derivative doped with ≥ 1 anthracene derivative and/or coumarin derivative and/or an electron injecting/transporting layer comprising a diphenylanthracene derivative with benzazole derivs. attached to the Ph groups. Application to displays is indicated.

L6 ANSWER 4 OF 37 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Organic electroluminescent device and novel thiophene derivative

Author/Inventor

Ishida, Tsutomu; Shimamura, Takehiko; Tanabe, Yoshimitsu; Totani, Yoshiyuki; Nakatsuka, Masakatsu

Patent Assignee/Corporate Source

Mitsui Chemicals Inc., Japan

Source

Jpn. Kokai Tokkyo Koho, 48 pp. CODEN: JKXXAF

Document Type

Patent

Language

Japanese

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
------------	------	------	-----------------	------

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2003282268	A	20031003	JP 2002-112966	20020416

Patent Number (1)
JP 2003282268
Kind Code (1)
A
Patent Publication Date (1)
20031003
Application Number (1)
JP 2002-112966
Application Date (1)
20020416
Priority Patent Number (1)
JP 2002-9104
Priority Kind Code (1)
A
Priority Patent Publication Date (1)
20020117

Abstract

The invention refers to an organic electroluminescent device comprising a novel thiophene derivative I [Ar = (un)substituted anthryl; Z1-3 = H, halo, straight chain, branched or cyclic alkyl, alkoxy, (un)substituted amino, aryl or aralkyl] in at least one layer.

L6 ANSWER 5 OF 37 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Anthracene compounds, their organic EL device materials, and their EL devices having high emission efficiency, long service life, and good heat resistance

Author/Inventor

Hosokawa, Chishio; Funabashi, Masakazu; Ikeda, Shuji; Yamamoto, Hiroshi

Patent Assignee/Corporate Source

Idemitsu Kosan Co., Ltd., Japan

Source

Jpn. Kokai Tokkyo Koho, 23 pp. CODEN: JKXXAF

Document Type

Patent

Language

Japanese

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2003238534	A	20030827	JP 2002-45705	20020222

Patent Number (1)
JP 2003238534
Kind Code (1)
A
Patent Publication Date (1)
20030827
Application Number (1)
JP 2002-45705
Application Date (1)
20020222
Priority Patent Number (1)
JP 2002-45705
Priority Patent Publication Date (1)
20020222

Abstract

The anthracene compds. are represented by a general formula of I [R1-R4, R6-R9 = H, halo, OH, NO₂, CN, amino, C1-30 alkyl, C4-40 alkenyl, CO₂H, etc.; R5 = divalent or trivalent C5-40 aromatic, divalent or trivalent C2-40 aromatic heterocyclic; R1-R9 may be bonded to neighboring group and form ring; A, B = C6-40 aryl, aromatic C2-40 heterocyclic; when R5 = C10-40 aromatic or aromatic C5-40 heterocyclic, A may be H; Ar1, Ar2 = C6-40 aryl, aromatic C2-40 heterocyclic, may be bonded to each other via linkage group L; L = (CR10R11)m, (SiR10R11)m, NR12m, vinylene, C6-40 arylene; R10-R12 = H, halo, C1-40 alkyl, C5-40 cycloalkyl, C5-40 aromatic hydrocarbyl, aromatic C2-40 heterocyclic, C7-40 aralkyl; m = 1, 2, 3; n = 0, 1]. The organic EL device contains, between anodes and cathodes, ≥1 organic thin-film layers involving a luminescent layer and containing I in ≥1 of the layers. Preferably, the organic thin-film layers consist of a luminescent layer, an electron-transporting layer, and a hole-transporting layer and at least the luminescent layer contains I. Preferably, the luminescent layer further contains arylamine compds. which may be selected from those represented by a general formula of Ar5(NAr6Ar7)p (Ar5 = C6-40 aromatic; Ar6, Ar7 = H, C6-40 aromatic; p = 1-6 integer) or Ar8(NAr9)qAr10rAr11Ar12s(NAr13)tAr14 (Ar8, Ar14 = C6-40 aromatic; Ar9-Ar13 = H, C6-40 aromatic; q, r, s, t = 0, 1). The electron-transporting layer may contain inorg. compds., preferably selected from dielects., semiconductors, or fine-crystalline or amorphous dielec. thin films. The dielects. may comprise ≥1 compds. selected from alkali metal chalcogenides, alkaline earth metal chalcogenides, alkali metal halides, and alkaline earth metal halides. The semiconductors may comprise ≥1 oxides, nitrides, or oxynitrides of ≥1 elements selected from Ba, Ca, Sr, Yb, Al, Ga, In, Li, Na, Cd, Mg, Si, Ta, Sb, and Zn. The electron-transporting layer may contain reducing dopants, preferably, ≥1 alkali metals selected from Na, K, Rb, and Cs and/or ≥1 alkaline earth metals

selected from Ca, r, and/or Ba. In another alternative, the organic thin-film layers consist of an electron-transporting layer, and a hole-transporting layer and at least one of these layers contain I.

L6 ANSWER 6 OF 37 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Organic electroluminescent device comprising coupled anthracene fluorene derivative and with amino-substituted hydrocarbon

Author/Inventor

Totani, Yoshiyuki; Ishida, Tsutomu; Shimamura, Takehiko; Tanabe, Yoshimitsu; Nakatsuka, Masakatsu

Patent Assignee/Corporate Source

Mitsui Chemicals Inc., Japan

Source

Jpn. Kokai Tokkyo Koho, 122 pp. CODEN: JKXXAF

Document Type

Patent

Language

Japanese

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2003229273	A	20030815	JP 2002-25736	20020201

Patent Number (1)

JP 2003229273

Kind Code (1)

A

Patent Publication Date (1)

20030815

Application Number (1)

JP 2002-25736

Application Date (1)

20020201

Priority Patent Number (1)

JP 2002-25736

Priority Patent Publication Date (1)

20020201

Abstract

The invention refers to an organic electroluminescent device comprising one or two fluorene rings directed bonded to an anthracene and a amino-substituted hydrocarbon.

L6 ANSWER 7 OF 37 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Organic electroluminescent devices of high brightness and luminescent efficiency and anthracene derivatives therefor

Author/Inventor

Ishida, Tsutomu; Shimamura, Takehiko; Tanabe, Yoshimitsu; Totani, Yoshiyuki; Nakatsuka, Masakatsu

Patent Assignee/Corporate Source

Mitsui Chemicals Inc., Japan

Source

Jpn. Kokai Tokkyo Koho, 99 pp. CODEN: JKXXAF

Document Type

Patent

Language

Japanese

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2003128651	A	20030508	JP 2001-317783	20011016

Patent Number (1)

JP 2003128651

Kind Code (1)

A

Patent Publication Date (1)

20030508

Application Number (1)

JP 2001-317783

Application Date (1)

20011016

Priority Patent Number (1)

JP 2001-317783

Priority Patent Publication Date (1)

20011016

Abstract

The anthracene derivs. have direct bonds between anthracene ring and fluorene ring and bear group I (Ar1, Ar2 = arylene; Z = bridging group).

L6 ANSWER 8 OF 37 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Organic electroluminescent device using pyrazole or pyrazoline

Author/Inventor

Suzuki, Koichi; Ueno, Kazunori; Senoo, Akihiro

Patent Assignee/Corporate Source

Canon Inc., Japan

Source

Jpn. Kokai Tokkyo Koho, 25 pp. CODEN: JKXXAF

Document Type

Patent

Language

Japanese

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2003109765	A	20030411	JP 2001-300548	20010928

Patent Number (1)

JP 2003109765

Kind Code (1)

A

Patent Publication Date (1)

20030411

Application Number (1)

JP 2001-300548

Application Date (1)

20010928

Priority Patent Number (1)

JP 2001-300548

Priority Patent Publication Date (1)

20010928

Abstract

The invention refers to an organic electroluminescent device comprising pyrazole or pyrazoline I [R1 = H, alkyl, (un)substituted aralkyl, aryl, heterocyclic, condensed polycyclic aromatic or heterocyclic; Ar1-3 = (un)substituted aryl, heterocyclic, condensed polycyclic aromatic or heterocyclic; at least two of R1, Ar1-3 are (un)substituted condensed polycyclic aromatic or heterocyclic groups].

L6 ANSWER 9 OF 37 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Organic electroluminescent devices

Author/Inventor

Suzuki, Koichi

Patent Assignee/Corporate Source

Canon Inc., Japan

Source

Jpn. Kokai Tokkyo Koho, 26 pp. CODEN: JKXXAF

Document Type

Patent

Language

Japanese

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2003109763	A	20030411	JP 2001-300546	20010928

Patent Number (1)

JP 2003109763

Kind Code (1)

A

Patent Publication Date (1)

20030411

Application Number (1)

JP 2001-300546

Application Date (1)

20010928

Priority Patent Number (1)

JP 2001-300546

Priority Patent Publication Date (1)

20010928

Abstract

The devices comprise a phosphor layer comprising R1-4Ar1, where R1-4 = H, alkyl, (substituted) aralkyl, (substituted) aryl, (substituted) heterocyclic, (substituted) condensed polyarom. ring, (substituted) polyheterocyclic ring, Ar1 = divalency-tetravalency

naphthylene, fluorenylene, anthracenylen, phenantrenylene, vinylene, triphenylene, thiophenylene, pyridylene, pyradylene, pyrimidiene, pyradylene, pyrimidiene, pyradadiene.

L6 ANSWER 10 OF 37 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Organic electroluminescent device and blue luminescence component

Author/Inventor

Sato, Hideki; Sato, Yoshiharu; Ichinosawa, Akiko

Patent Assignee/Corporate Source

Mitsubishi Chemical Corp., Japan

Source

Jpn. Kokai Tokkyo Koho, 23 pp. CODEN: JKXXAF

Document Type

Patent

Language

Japanese

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2003031371	A	20030131	JP 2001-216944	20010717

Patent Number (1)

JP 2003031371

Kind Code (1)

A

Patent Publication Date (1)

20030131

Application Number (1)

JP 2001-216944

Application Date (1)

20010717

Priority Patent Number (1)

JP 2001-216944

Priority Patent Publication Date (1)

20010717

Abstract

The invention refers to an electroluminescent device comprising I [Z = divalent substituent; and the Ph and carbazoyl groups may be substituted] as a hole blocking layer.

L6 ANSWER 11 OF 37 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Organic electroluminescent device and luminescent material containing a rubrene derivative as a luminescence facilitator

Author/Inventor

Hamada, Yuji; Tsujioka, Tsuyoshi

Patent Assignee/Corporate Source

Sanyo Electric Co., Ltd., Japan

Source

PCT Int. Appl., 176 pp. CODEN: PIXXD2

Document Type

Patent

Language

Japanese

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002100977	A1	20021219	WO 2002-JP5522	20020604

Patent Number (1)

WO 2002100977

Kind Code (1)

A1

Patent Publication Date (1)

20021219

Application Number (1)

WO 2002-JP5522

Application Date (1)

20020604

Priority Patent Number (1)

JP 2001-171664

Priority Kind Code (1)

A

Priority Patent Publication Date (1)

20010606

Abstract

An organic electroluminescence device having a high luminance and a high emission efficiency. The invention refers to an organic electroluminescent device comprising a luminescent layer with a host material, an emitting dopant and a luminescence facilitator, wherein the facilitator is a rubrene derivative

L6 ANSWER 12 OF 37 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Electroluminescence Characteristics of Card Anthracene-containing Polyimide: The Effect of the Cathode and Anode Materials

Author/Inventor

Kolesnikov, V. A.; Brusentseva, M. A.; Rumyantsev, B. M.; Berendyaev, V. I.; Vannikov, A. V.

Patent Assignee/Corporate Source

Frumkin Institute of Electrochemistry, Russian Academy of Sciences, Moscow, 119071, Russia

Source

Russian Journal of Electrochemistry (Translation of Elektrokhimiya) (2002), 38(11), 1163-1172 CODEN: RJELE3; ISSN: 1023-1935

Document Type

Journal

Language

English

Abstract

The effect of electrode materials on the characteristics of electroluminescence devices of the type transparent conductive substrate/polyimide/metallic electrode is studied. ITO. The reasons for variations in the electroluminescence spectra after replacing the substrate material and the metal of the top electrode are discussed. The applicability of the Fauler-Nordheim model for describing the injection of charge carriers in the electroluminescence devices under study is analyzed.

L6 ANSWER 13 OF 37 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Double-spiro organic compounds and electroluminescent devices

Author/Inventor

Kim, Kong-Kyeum; Son, Se-Hwan; Yoon, Seok-Hee; Bae, Jae-Soon; Lee, Youn-Gu; Im, Sung-Gap; Kim, Ji-Eun; Lee, Jae-Chol

Patent Assignee/Corporate Source

LG Chem, Ltd., S. Korea

Source

PCT Int. Appl., 117 pp. CODEN: PIXXD2

Document Type

Patent

Language

English

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002088274	A1	20021107	WO 2002-KR458	20020318

Patent Number (1)

WO 2002088274

Kind Code (1)

A1

Patent Publication Date (1)

20021107

Application Number (1)

WO 2002-KR458

Application Date (1)

20020318

Priority Patent Number (1)

KR 2001-23038

Priority Kind Code (1)

A

Priority Patent Publication Date (1)

20010427

Abstract

Double-spiro organic compds. are claimed which are described by the general formula I (R1-24 = independently selected substituents not all of which are H). Light-emitting, hole-transporting, and electron-transporting materials comprising the compds. are also described; Electroluminescent materials comprising the compds, including deposited films, methods for depositing the materials, and organic electroluminescent devices employing the materials, and method for fabricating the devices, are also described.

L6 ANSWER 14 OF 37 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Near-infrared electroluminescence in polymer composites based on organic nanocrystals

Author/Inventor

Maltsev, Eugene I.; Lypenko, Dmitry A.; Bobinkin, Vladimir V.; Tameev, Alek R.; Kirillov, Sergey V.; Shapiro, Boris I.; Schoo, Herman F. M.; Vannikov, Anatoly V.

Patent Assignee/Corporate Source

Frumkin Institute of Electrochemistry of the Russian Academy of Sciences, Moscow, 117071, Russia

Source Applied Physics Letters (2002), 81(16), 3088-3090 CODEN: APPLAB; ISSN: 0003-6951
Document Type Journal
Language English

Abstract IR electroluminescence was revealed in single-layer light-emitting diodes based on a type of electroactive polymer nanocomposites-electron-hole conducting aromatic polyimide and organic nanocryst. particles of cyanine mols., known as J-aggregates. These materials exhibit a very narrow emission band with a maximum at 815 nm. Dramatic increase of charge-carrier mobility was observed for these layers containing the J-aggregate nanocryst. phase.

L6 ANSWER 15 OF 37 CAPLUS COPYRIGHT 2008 ACS on STN
Title

Electroluminescent (EL) devices

Author/Inventor Hu, Nan-Xing; Aziz, Hany; Jain, Poonam; Popovic, Zoran D.
Patent Assignee/Corporate Source Xerox Corporation, USA

Source U.S. Pat. Appl. Publ., 46 pp. CODEN: USXXCO
Document Type Patent
Language English

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2002132134	A1	20020919	US 2001-771311	20010126

Patent Number (1)
US 2002132134
Kind Code (1)
A1

Patent Publication Date (1)
20020919
Application Number (1)
US 2001-771311

Application Date (1)
20010126
Priority Patent Number (1)
US 2001-771311

Priority Kind Code (1)
A3
Priority Patent Publication Date (1)
20010126

Abstract Electroluminescent devices are described which employ compds. are described by the general formula I and II (R1 and R2 = H, alkyl, alicyclic alkyl, alkoxy, halo, and cyano groups, and, in II, aryl groups; Ar1 and Ar2 = independently selected aromatic component or an aryl group comprised of 4-15 conjugate-bonded or fused benzene rings; R3, R4, R5, and R6 = independently selected H, an alkyl, alicyclic alkyl, aryl, and alkoxy group; wherein R3 and R4, or R4 and R5 are optionally combined into a bivalent hydrocarbon group selected from the group consisting of an alkylene, an alkylidene, an alicyclic alkylidene, and an arylalkylidene; Ar3 and Ar4 = independently selected aryl groups; and Ar = a tetravalent aromatic group). The compds. and their mixts. are also described.

L6 ANSWER 16 OF 37 CAPLUS COPYRIGHT 2008 ACS on STN
Title

Spiro compound for organic electroluminescent material and component

Author/Inventor Enokida, Toshio; Toba, Yasumasa
Patent Assignee/Corporate Source Toyo Ink Mfg. Co., Ltd., Japan

Source Jpn. Kokai Tokkyo Koho, 48 pp. CODEN: JKXXAF
Document Type Patent
Language Japanese

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2002265938	A	20020918	JP 2001-72095	20010314

Patent Number (1)

JP 2002265938

Kind Code (1)

A

Patent Publication Date (1)

20020918

Application Number (1)

JP 2001-72095

Application Date (1)

20010314

Priority Patent Number (1)

JP 2001-72095

Priority Patent Publication Date (1)

20010314

Abstract

The invention refers to an electroluminescent component comprising the spiro compound I [R1-16, X1,2 = H, halo, cyano, nitro, (un)substituted alkyl, alkoxy, aryloxy alkylthio, arylthio, siloxy, cycloalkyl, aryl, aromatic heterocycle, amino, styryl and adjacent groups or their substituents may join to form rings] as a luminescent material.

L6 ANSWER 17 OF 37 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Metal coordination compound, luminescence device and display apparatus

Author/Inventor

Takiguchi, Takao; Okada, Shinjiro; Tsuboyama, Akira; Miura, Seishi; Moriyama, Takashi; Kamatani, Jun; Furugori, Manabu

Patent Assignee/Corporate Source

Canon Kabushiki Kaisha, Japan

Source

Eur. Pat. Appl., 113 pp. CODEN: EPXXDW

Document Type

Patent

Language

English

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 1238981	A2	20020911	EP 2002-5112	20020307

Patent Number (1)

EP 1238981

Kind Code (1)

A2

Patent Publication Date (1)

20020911

Application Number (1)

EP 2002-5112

Application Date (1)

20020307

Priority Patent Number (1)

JP 2001-64254

Priority Kind Code (1)

A

Priority Patent Publication Date (1)

20010308

Abstract

An electroluminescence device having a layer containing a specific metal coordination compound is provided. The metal coordination compound is represented by $ML_mL'_n$ ($M = Ir, Pt, Rh$ or Pd ; L and L' = mutually different bidentate ligands; $m = 1-3$; $n = 0-2$; $m+n = 2$ or 3 ; a partial structure ML_m is represented by I; a partial structure ML'_n is represented by II (CyN1 and CyN2 = cyclic group capable of having a substituent, including a nitrogen and bonded to the metal atom M via the nitrogen atom; CyC1 and CyC2 = cyclic group capable of having a substituent selected from a halogen atom, cyano group, nitro, trialkylsilyl, etc.; the cyclic groups CyC1 and CyC2 include an aromatic group capable of having a substituent represented by IV ($Y = CO, C=C(CN)_2, O, S, CRR'$; $R, R' = H, C1-8$ alkyl, etc.)), and III ($E, G = C1-20$ alkyl of which the H can be optionally replaced with a F, or an aromatic group capable of having a substituent)). The metal coordination compound having the aromatic group is effective in providing high-efficiency luminescence and long-term high luminance.

L6 ANSWER 18 OF 37 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Blue-Emitting Anthracenes with End-Capping Diarylamines

Author/Inventor

Danel, Krzysztof; Huang, Tai-Hsiang; Lin, Jiann T.; Tao, Yu-Tai; Chuen, Chang-Hao

Patent Assignee/Corporate Source

Institute of Chemistry, Academia Sinica, Taipei, WA, 115, USA

Source

Chemistry of Materials (2002), 14(9), 3860-3865 CODEN: CMATEX; ISSN: 0897-4756

Document Type

Journal
Language English

Abstract 2-Tert-butyl-9,10-bis(bromoaryl)anthracenes were synthesized from 2-tert-butyl-9,10-anthraquinone. Pd-catalyzed C-N bond formation between these bromo compds. and diarylamines provides stable 2-tert-butyl-9,10- diarylanthracenes containing two peripheral diarylamines (anth). They possess high thermal decomposition temperature ($T_d > 450^\circ$) and form a stable glass ($T_g > 130^\circ$), also, they are fluorescent in the blue region with moderate to good quantum efficiencies. Two types of light-emitting diodes (LED) were constructed from anth, (I) ITO/anth/TPBI/Mg:Ag and (II) ITO/anth/Alq3/Mg:Ag, where TPBI and Alq3 are 1,3,5-tris(N-phenylbenzimidazol-2-yl)benzene and tris(8-hydroxyquinolato)aluminum, resp. In type I devices, the anth function as the hole-transporting and emitting material. In type II devices, emission from Alq3 is observed. Several blue-light-emitting type I devices exhibit good maximum brightness and phys. performance. The relation between the energy levels of the anth and the performance of the light-emitting diode is discussed.

L6 ANSWER 19 OF 37 CAPLUS COPYRIGHT 2008 ACS on STN

Title Organic polymeric light-emitting devices
Author/Inventor Vannikov, A. V.
Patent Assignee/Corporate Source Inst. Elektrokhim. im. A. N. Frumkina, RAN, Moscow, 117071, Russia

Source Rossiiskii Khimicheskii Zhurnal (2001), 45(5-6), 41-50 CODEN: RKZHEZ; ISSN: 1024-6215

Document Type Journal

Language Russian

Abstract The author developed new materials for polymeric light-emitting devices. The preparation methods can be used in other optoelectronics. The electronic transport in polymer layers is discussed.

L6 ANSWER 20 OF 37 CAPLUS COPYRIGHT 2008 ACS on STN

Title The organic electroluminescent material and organic electroluminescent component

Author/Inventor Suda, Yasumasa; Onikubo, Shunichi
Patent Assignee/Corporate Source Toyo Ink Mfg. Co., Ltd., Japan

Source Jpn. Kokai Tokkyo Koho, 33 pp. CODEN: JKXXAF

Document Type Patent

Language Japanese

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2002060742	A	20020226	JP 2000-243895	20000811

Patent Number (1)
JP 2002060742

Kind Code (1)
A

Patent Publication Date (1)
20020226

Application Number (1)
JP 2000-243895

Application Date (1)
20000811

Priority Patent Number (1)
JP 2000-243895

Priority Patent Publication Date (1)
20000811

Abstract The invention refers to an organic electroluminescent material comprising R1R2C.CR3R4-R5 [R1-3 = aryl, heterocyclic; R4 = divalent condensed polycyclic hydrocarbon or divalent heterocyclic R5 = H, unsubstituted aryl, heterocyclic, or a combination thereof].

L6 ANSWER 21 OF 37 CAPLUS COPYRIGHT 2008 ACS on STN

Title Hydrocarbon compound for organic electroluminescent elements and using them
Author/Inventor

Ishida, Tsutomu; Shimamura, Takehiko; Totani, Yoshiyuki; Nakatsuka, Masakatsu
Patent Assignee/Corporate Source
Mitsui Chemicals, Inc., Japan

Source PCT Int. Appl., 251 pp. CODEN: PIXXD2

Document Type
Patent

Language
Japanese

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002014244	A1	20020221	WO 2001-JP6920	20010810

Patent Number (1)
WO 2002014244

Kind Code (1)
A1

Patent Publication Date (1)
20020221

Application Number (1)
WO 2001-JP6920

Application Date (1)
20010810

Priority Patent Number (1)
JP 2000-242476

Priority Kind Code (1)
A

Priority Patent Publication Date (1)
20000810

Abstract

Title electroluminescent elements comprise one pair of electrodes and pinched between the electrodes, ≥ 1 layer(s) containing ≥ 1 novel hydrocarbon compound in a general formula $X1(F1)(A1)k(F2)(A2)m(F3)nX2$ [A1-2 = (un)substituted anthracenediyl; F1-3 = (un)substituted fluorenediyl; X1-2 = H, halo, straight, branched or cyclic alkyl, alkoxy, amino, aryl, or (un)substituted amino, aryl or aralkyl, j,m,n = 0, 1; k,l = 1, 2] having an anthracene ring and a fluorene ring which are directly bonded with each other. The compound can be suitably used for preparing an organic electroluminescent element being excellent in luminous efficiency and having a long luminous life.

L6 ANSWER 22 OF 37 CAPLUS COPYRIGHT 2008 ACS on STN

Title New heterocyclic compound for electroluminescent device
Author/Inventor

Okada, Hisashi; Ise, Toshihiro
Patent Assignee/Corporate Source
Fuji Photo Film Co., Ltd., Japan

Source Jpn. Kokai Tokkyo Koho, 52 pp. CODEN: JKXXAF

Document Type
Patent

Language
Japanese

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2001335776	A	20011204	JP 2000-218967	20000719

Patent Number (1)
JP 2001335776

Kind Code (1)
A

Patent Publication Date (1)
20011204

Application Number (1)
JP 2000-218967

Application Date (1)
20000719

Priority Patent Number (1)
JP 1999-207957

Priority Kind Code (1)
A

Priority Patent Publication Date (1)
19990722

Abstract

The invention relates to new heterocyclic compds., suited for use in making an electroluminescent device, represented by L-(A)m [A

= heterocyclic group having ≥ 2 aromatic hetero ring condensed; m = integer ≥ 2 ; L = bonding group].

L6 ANSWER 23 OF 37 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Organic electroluminescent device

Author/Inventor

Kitazawa, Daisuke; Tominaga, Takeshi; Kohama, Toru

Patent Assignee/Corporate Source

Toray Industries, Inc., Japan

Source

Jpn. Kokai Tokkyo Koho, 20 pp. CODEN: JKXXAF

Document Type

Patent

Language

Japanese

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2001223082	A	20010817	JP 2000-30373	20000208

Patent Number (1)

JP 2001223082

Kind Code (1)

A

Patent Publication Date (1)

20010817

Application Number (1)

JP 2000-30373

Application Date (1)

20000208

Priority Patent Number (1)

JP 2000-30373

Priority Patent Publication Date (1)

20000208

Abstract

The invention relates to an organic electroluminescent device comprising the compound represented by I [R1-R14 = H, alkyl, cycloalkyl, etc.; at least one of R3 and R4 have steric hindrance to limit the free internal rotation] and a 580-720 nm emitting fluorescent material having a pyrromethane skeleton.

L6 ANSWER 24 OF 37 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Charge carrier transport in aromatic polyimides and polyimide/J-aggregate composites

Author/Inventor

Tameev, Alek R.; Kozlov, Aleksey A.; Mal'tsev, Eugene I.; Lypenko, Dmitry A.; Bobonkin, Vladimir V.; Vannikov, Anatoly V.

Patent Assignee/Corporate Source

Frumkin Institute of Electrochemistry of the Russian Academy of Sciences, Moscow, 117071, Russia

Source

Proceedings of SPIE-The International Society for Optical Engineering (2001), 4105(Organic Light-Emitting Materials and Devices IV), 443-449 CODEN: PSISDG; ISSN: 0277-786X

Document Type

Journal

Language

English

Abstract

Charge carrier transport in aromatic polyimides based on 9,10-bis(aminophenyl)anthracene or 9,10-bis(phenylthio)anthracene and their composites with dye J-aggregates was studied using conventional time-of-flight techniques. The elec. field and temperature dependencies of both hole and electron drift mobility were observed. In amorphous films of the soluble polyimide, the drift mobility was found to reach the value of 10-4 cm² V⁻¹ s⁻¹. In films of the insol. polyimide containing a crystalline phase, the mobility was lower by one or two orders of magnitude. The result is attributed to the presence of cavities in the crystalline film. The applicability of known theor. models describing the temperature and elec. field dependencies of mobility is discussed. J-aggregates, formed in the soluble polyimides doped with cyanine dye mols., play an active role in charge carrier transport in the electroluminescence composites.

L6 ANSWER 25 OF 37 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Electroluminescence component

Author/Inventor

Tanaka, Hiromitsu; Mouri, Makoto; Takeuchi, Hisato; Tokito, Seishi

Patent Assignee/Corporate Source

Toyota Central Research and Development Laboratories, Inc., Japan

Source

Jpn. Kokai Tokkyo Koho, 32 pp. CODEN: JKXXAF

Document Type

Patent

Language

Japanese

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2001110572	A	20010420	JP 2000-237442	20000804

Patent Number (1)

JP 2001110572

Kind Code (1)

A

Patent Publication Date (1)

20010420

Application Number (1)

JP 2000-237442

Application Date (1)

20000804

Priority Patent Number (1)

JP 1999-221653

Priority Kind Code (1)

A

Priority Patent Publication Date (1)

19990804

Abstract

The invention refers to an electroluminescent device comprising two electrodes and an electroluminescent layer containing I [A1,2 = functional group; B1-6 = direct bonds or divalent functional groups; A1,2 = triphenylamine, coumarin, or oxadiazole derivative groups having hole and electron transport and luminescent properties].

L6 ANSWER 26 OF 37 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Organic electroluminescence and organic luminous medium

Author/Inventor

Hosokawa, Chishio; Higashi, Hisahiro; Fukuoka, Kenichi; Ikeda, Hidetsugu

Patent Assignee/Corporate Source

Idemitsu Kosan Co., Ltd., Japan

Source

PCT Int. Appl., 41 pp. CODEN: PIXXD2

Document Type

Patent

Language

Japanese

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001021729	A1	20010329	WO 2000-JP6402	20000920

Patent Number (1)

WO 2001021729

Kind Code (1)

A1

Patent Publication Date (1)

20010329

Application Number (1)

WO 2000-JP6402

Application Date (1)

20000920

Priority Patent Number (1)

JP 1999-267460

Priority Kind Code (1)

A

Priority Patent Publication Date (1)

19990921

Abstract

The invention refers to a organic electroluminescent device comprising a mono-, di- or tri- styryl amine, and at least one of the anthracene derivs., A1LA1 [A1,2 = (un)substituted mono Ph anthryl, or (un)substituted di-Ph anthryl; L = single bond or divalent chain] and A3AnA4 [An = (un)substituted anthracene; A3,4 = (un)substituted condensed aromatic ring, or (un)substituted C12+ chain uncondensed aryl ring].

L6 ANSWER 27 OF 37 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Organic electroluminescent devices

Author/Inventor

Sato, Tadahisa; Hara, Shintaro
Patent Assignee/Corporate Source
Fuji Photo Film Co., Ltd., Japan; Matsushita Electric Industrial Co., Ltd.

Source
Jpn. Kokai Tokkyo Koho, 24 pp. CODEN: JKXXAF

Document Type
Patent

Language
Japanese

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2000323281	A	20001124	JP 1999-135920	19990517

Patent Number (1)
JP 2000323281

Kind Code (1)
A

Patent Publication Date (1)
20001124

Application Number (1)
JP 1999-135920

Application Date (1)
19990517

Priority Patent Number (1)
JP 1999-135920

Priority Patent Publication Date (1)
19990517

Abstract

The devices comprise a hole transport layer comprising I, II, III, IV or V (A1-9, B1-9, C1-9 = (substituted) ethylene, (substituted) vinylene, (substituted) o-arylene; Ar1-5 = (substituted) aromatic hydrocarbon, (substituted) aromatic heterocyclic hydrocarbon; a, b, c = 1-4; d = 0-2; Ar6-8 = Ar1-5 when Y = N; Ar6-8 = (substituted) benzene ring when Y = 1,3,5-benzenetolyl; e, f, g = 1-3; Ar9 = Ar1-5 except benzene ring, (substituted) polyaryl methane; h = 1-4; Ar10,11 = Ar1-5; i, k = 1-4; j ≥ 1; Z = 1-4 valent group of aromatic ring, aromatic heterocyclic, triarylamine, polyarylethane; m = 1-4; l ≥ 1; n = 1-4).

L6 ANSWER 28 OF 37 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Electroluminescent properties of anthracene-containing polyimides

Author/Inventor

Mal'tsev, Eugene I.; Brusentseva, Maria A.; Lypenko, Dmitry A.; Berendyaev, Vladimir I.; Kolesnikov, Vladislav A.; Kotov, Boris V.; Vannikov, Anatoly V.

Patent Assignee/Corporate Source

Frumkin Institute of Electrochemistry of the Russian Academy of Sciences, Moscow, 117071, Russia

Source

Polymers for Advanced Technologies (2000), 11(7), 325-329 CODEN: PADTE5; ISSN: 1042-7147

Document Type

Journal

Language

English

Abstract

Optical and electroluminescent properties of a new soluble anthracene-containing polyimide (ACPI) was studied. Solubility of ACPI in organic solvents allows direct spin casting of the polymer films exhibiting intense photo- and electroluminescence (EL) in the visible range. This nonconjugated polymer was used as emitting and electron-hole transporting layers in polymer light-emitting devices (LEDs). EL properties of the uni- and bilayer LEDs are discussed in terms of the band structure, bipolar transport and electron donor-acceptor interactions.

L6 ANSWER 29 OF 37 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Aromatic hydrocarbon compound for organic electroluminescent device

Author/Inventor

Funabashi, Masakazu; Kawamura, Hisayuki; Azuma, Hisaohiro; Hosokawa, Chishio

Patent Assignee/Corporate Source

Idemitsu Kosan Co., Ltd., Japan

Source

Jpn. Kokai Tokkyo Koho, 18 pp. CODEN: JKXXAF

Document Type

Patent

Language

Japanese

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2000192028	A	20000711	JP 1998-370480	19981225

Patent Number (1)
JP 2000192028
Kind Code (1)
A
Patent Publication Date (1)
20000711
Application Number (1)
JP 1998-370480
Application Date (1)
19981225
Priority Patent Number (1)
JP 1998-370480
Priority Patent Publication Date (1)
19981225

Abstract

An aromatic hydrocarbon compound for an organic electroluminescent device is represented by I [Ar1 and Ar2 = arylene and heterocyclic groups; Ar3 and Ar4 = aryl and heterocyclic groups; R represents a group combined with either Ar2 or Ar3 to form a ring; n = 0-2 integer; and m = 0 or 1].

L6 ANSWER 30 OF 37 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Electroluminescent properties of anthracene-containing polyimides

Author/Inventor

Mal'tsev, Eugene I.; Brusentseva, Mariy A.; Berendyaev, Vladimir I.; Kolesnikov, Vladislav A.; Kotov, Boris V.; Vannikov, Anatoly V.

Patent Assignee/Corporate Source

Frumkin Institute of Electrochemistry, Russian Academy of Sciences, Moscow, Russia

Source

Proceedings of SPIE-The International Society for Optical Engineering (1999), 3797(Organic Light-Emitting Materials and Devices II), 350-358 CODEN: PSISDG; ISSN: 0277-786X

Document Type

Journal

Language

English

Abstract

The electroluminescence (EL) of donor-acceptor polyimides prepared from 9,10-bis(m-aminophenylthio)-anthracene (BPTA) and 1,3-bis(3,4-dicarboxyphenoxy)benzene or 2,2-bis[4-(3,4-dicarboxyphenoxy)phenyl]propane dianhydrides was studied. The aromatic polyimides with and without sulfur atoms in the backbone, were evaluated as electron-hole transporting and light-emitting materials for use in single- and multilayer electroluminescent diodes. These polyimides are efficient electron and hole conductors and also exhibit intense photoluminescence of exciplex origin. Some of the polyimides have been used as hole conducting layers with tris(8-quinolinolato)aluminum complex (Alq3) as electron conducting layer in bilayer LEDs of high brightness. A direct correlation was revealed between transport characteristics and electroluminescent properties of these electroactive materials. At room temperature, the electron mobility and hole drift mobility directly measured by conventional TOF techniques indicate effective bipolar transport. The simplicity of synthesis, high thermal stability, organic solvent solubility, and excellent film-forming ability make these polyimides good candidates for technol. applications. The band structure, bipolar transport, and electron donor-acceptor interactions in test one-layer and bilayer LEDs based on the polyimides are described.

L6 ANSWER 31 OF 37 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Organic electroluminescent devices

Author/Inventor

Hosokawa, Chishio; Funehashi, Masakazu; Kawamura, Hisayuki; Arai, Hiromasa; Koga, Hidetoshi; Ikeda, Hidetsugu

Patent Assignee/Corporate Source

Idemitsu Kosan Co., Ltd., Japan

Source

PCT Int. Appl., 167 pp. CODEN: PIXXD2

Document Type

Patent

Language

Japanese

Patent Information

PATENT NO	KIND	DATE	APPLICATION NO.	DATE
WO 2000039247	A1	20000706	WO 1999-JP7390	19991228

Patent Number (1)
WO 2000039247
Kind Code (1)
A1
Patent Publication Date (1)
20000706
Application Number (1)
WO 1999-JP7390

Application Date (1)
19991228
Priority Patent Number (1)
JP 1998-373921
Priority Kind Code (1)
A
Priority Patent Publication Date (1)
19981228

Abstract

The devices having a high luminescent efficiency, a long life and a high heat resistance comprise I (A = (substituted) C22-60 arylene; X1-4 = (substituted) C6-30 arylene; Y1-4 = II; a-d = 0-2; R1-4 = H, (substituted) alkyl, (substituted) aryl, cyano; R3 may be bonded to R4 to form a triple bond; Z = (substituted) aryl; n = 0, 1).

L6 ANSWER 32 OF 37 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Aromatic hydrocarbon compound for organic electroluminescent device

Author/Inventor

Azuma, Hisahiro; Hosokawa, Chishio; Kusumoto, Tadashi

Patent Assignee/Corporate Source

Idemitsu Kosan Co., Ltd., Japan

Source

Jpn. Kokai Tokkyo Koho, 17 pp. CODEN: JKXXAF

Document Type

Patent

Language

Japanese

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2000053676	A	20000222	JP 1998-225679	19980810

Patent Number (1)

JP 2000053676

Kind Code (1)

A

Patent Publication Date (1)

20000222

Application Number (1)

JP 1998-225679

Application Date (1)

19980810

Priority Patent Number (1)

JP 1998-225679

Priority Patent Publication Date (1)

19980810

Abstract

The aromatic hydrocarbon compound for organic electroluminescent device has structure (R1)(R2)C=CH-Ar1-An-Ar2-CH=C(R3)(R4) (An = C1-30 alkyl, alkoxy, C6-18 aryloxy, amino, etc.; Ar1-2 = divalent heterocyclic ring with C5-30 and S, polyarylene; R1-4 = H, C1-30 alkyl, alkoxy, C6-30 aryl, etc.). The aromatic hydrocarbon compound provides the organic electroluminescent device of the high electroluminescent efficiency, the decreased driving voltage, and the excellent heat-resistance.

L6 ANSWER 33 OF 37 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Organic electroluminescent material containing anthracene derivative and organic electroluminescent device with it

Author/Inventor

Tamano, Michiko; Maki, Shinichiro; Onikubo, Shunichi; Okutsu, Satoshi; Enokida, Toshio

Patent Assignee/Corporate Source

Toyo Ink Mfg. Co., Ltd., Japan

Source

Jpn. Kokai Tokkyo Koho, 22 pp. CODEN: JKXXAF

Document Type

Patent

Language

Japanese

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 11111458	A	19990423	JP 1997-264468	19970929

Patent Number (1)

JP 11111458

Kind Code (1)

A

Patent Publication Date (1)
19990423
Application Number (1)
JP 1997-264468
Application Date (1)
19970929
Priority Patent Number (1)
JP 1997-264468
Priority Patent Publication Date (1)
19970929

Abstract

The material comprises an anthracene derivative having a formula I (A1, 2 = alkyl, alkoxy, aryloxy, condensed polycyclic, alkylamino, arylamino; R1-16 = H, halogen, cyano, NO₂, alkyl, alkoxy, aryloxy, alkylthio, arylthio, cyclic group, NH₂; R1-16 may bond to form a ring). The device has a light-emitting layer-containing plural organic compound thin films sandwiched between a pair of electrodes, at least one of the films contains the material. The device shows high luminance with efficiency and long life.

L6 ANSWER 34 OF 37 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Organic electroluminescent device material containing naphthalene derivative and organic electroluminescent device with it

Author/Inventor

Okutsu, Satoshi; Tamano, Michiko; Onikubo, Shunichi; Enokida, Toshio

Patent Assignee/Corporate Source

Toyo Ink Mfg. Co., Ltd., Japan

Source

Jpn. Kokai Tokkyo Koho, 28 pp. CODEN: JKXXAF

Document Type

Patent

Language

Japanese

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 10289786	A	19981027	JP 1997-95406	19970414

Patent Number (1)

JP 10289786

Kind Code (1)

A

Patent Publication Date (1)

19981027

Application Number (1)

JP 1997-95406

Application Date (1)

19970414

Priority Patent Number (1)

JP 1997-95406

Priority Patent Publication Date (1)

19970414

Abstract

The title material contains the derivative described by the general formula I (X = halo, cyano, alkyl, aryl, alkoxy, aryloxy, alkylthio, arylthio, cycloalkyl, heterocyclic, NH₂; i = 1-28). Device are also described which have plural organic compound thin films, containing a light-emitting layer and a hole injection layer, sandwiched by a pair of electrodes, in which one of the layers contains the material. The devices show high luminance, efficiency, and long life.

L6 ANSWER 35 OF 37 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Organic electroluminescent materials and devices using the same with high luminance and long life

Author/Inventor

Okutsu, Satoshi; Onikubo, Shunichi; Tamano, Michiko; Enokida, Toshio

Patent Assignee/Corporate Source

Toyo Ink Mfg. Co., Ltd., Japan

Source

Jpn. Kokai Tokkyo Koho, 20 pp. CODEN: JKXXAF

Document Type

Patent

Language

Japanese

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 10152676	A	19980609	JP 1996-313289	19961125

Patent Number (1)

JP 10152676

Kind Code (1)

A

Patent Publication Date (1)

19980609

Application Number (1)

JP 1996-313289

Application Date (1)

19961125

Priority Patent Number (1)

JP 1996-313289

Priority Patent Publication Date (1)

19961125

Abstract

Title materials are oxazole derivs. I [X1-3 = N, CH, or C bonding with Ar1 or Ar2, where X1 or X3 is C; Ar1-2 = arylene; Ar3-5 = H, cyano, (cyclo) alkyl, aryl, heterocycle; m, n = 0-4]. Electroluminescent devices including layers (preferably emitting layers) containing I are also claimed.

L6 ANSWER 36 OF 37 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Electroluminescence of anthracene-containing polyimides

Author/Inventor

Mat'nev, Evgenii I.; Brusentseva, Maria A.; Berendyaev, Vladimir I.; Kolesnikov, Vladislav A.; Lunina, Elena V.; Kotov, Boris V.; Vannikov, Anatolii V.

Patent Assignee/Corporate Source

A N Frumkin Institute of Electro-Chemistry, Russian Academy of Sciences, Moscow, 117071, Russia

Source

Mendeleev Communications (1998), (1), 31-32 CODEN: MENCEX; ISSN: 0959-9436

Document Type

Journal

Language

English

Abstract

Electroluminescence has been revealed in a new class of electroactive polymers, the anthracene-containing aromatic polyimide derivs.; high thermal stability, ability to cast layers from solution and excellent film-forming properties make these materials of potential interest for technol. applications.

L6 ANSWER 37 OF 37 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Light-emitting material for organo- electroluminescence device and organo- electroluminescence device for which the light-emitting material is adapted

Author/Inventor

Tamano, Michiko; Enokida, Toshio

Patent Assignee/Corporate Source

Toyo Ink Manufacturing Co., Ltd., Japan

Source

Eur. Pat. Appl., 31 pp. CODEN: EPXXDW

Document Type

Patent

Language

English

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 786926	A2	19970730	EP 1997-300551	19970129

Patent Number (1)

EP 786926

Kind Code (1)

A2

Patent Publication Date (1)

19970730

Application Number (1)

EP 1997-300551

Application Date (1)

19970129

Priority Patent Number (1)

JP 1996-12488

Priority Kind Code (1)

A

Priority Patent Publication Date (1)

19960129

Abstract

Comps. for use in electroluminescent devices are described by the general formulas I and II (A-D are the same or different groups

each = (un)substituted alkyl, (un)substituted monocyclic group, or (un)substituted fused polycyclic group, or A and B and/or C and D, together with the nitrogen atom to which they are attached, form a substituted or unsubstituted heterocyclic ring. R1-20 are independently selected from H, halogen atoms, (un)substituted alkyl, (un)substituted alkoxy, (un)substituted amino, (un)substituted monocyclic, or (un)substituted fused polycyclic groups; and X1-4 are independently selected from various linking groups). Television sets, light-emitting devices, copy machines, printers, liquid-crystal displays, displays, electrophotog. photoreceptors, photoelec. converters, solar cells, and image sensors containing electroluminescent devices employing the compds. are also described.

=>

Connecting via Winsock to STN

Welcome to STN International! Enter x:X



LOGINID:SSPTAMEN1774

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

=>



chain nodes :

12 13

ring nodes :

1 2 3 4 5 6 7 8 9

chain bonds :

8-12 12-13

ring bonds :

1-2 1-5 1-9 2-3 3-4 4-5 5-6 6-7 7-8 8-9

exact/norm bonds :

1-2 1-5 1-9 2-3 3-4 4-5 5-6 6-7 7-8 8-9 8-12 12-13

G1:C,N

Match level :

1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:Atom 8:Atom 9:Atom 12:Atom 13:Atom

L1 STRUCTURE UPLOADED

-> s 11 sss full
FULL SEARCH INITIATED 15:10:46 FILE 'REGISTRY'
FULL SCREEN SEARCH COMPLETED - 228780 TO ITERATE

100.0% PROCESSED 228780 ITERATIONS 4458 ANSWERS
SEARCH TIME: 00.00.03

L2 4458 SEA SSS FUL L1

=> s 12
L3 396 L2

=> s 13 and py<-2003
23975225 PY<-2003

L4 235 L3 AND PY<-2003

=> s 14 and (electroluminescent or electroluminescence)
68989 ELECTROLUMINESCENT
6 ELECTROLUMINESCENTS
68991 ELECTROLUMINESCENT
(ELECTROLUMINESCENT OR ELECTROLUMINESCENTS)
22386 ELECTROLUMINESCENCE
25 ELECTROLUMINESCENCES
22390 ELECTROLUMINESCENCE
(ELECTROLUMINESCENCE OR ELECTROLUMINESCENCES)

L5 1 L4 AND (ELECTROLUMINESCENT OR ELECTROLUMINESCENCE)

=> d 15 ibib abs

L5 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2008 ACS on STN

Title Organic electroluminescent device

Author/Inventor Nakatsuka, Masakatsu; Shimamura, Takehiko

Patent Assignee/Corporate Source Mitsui Chemicals Inc., Japan

Source Jpn. Kokai Tokkyo Koho, 43 pp. CODEN: JKXXAF

Document Type Patent

Language Japanese

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2001035664	A	20010209	JP 1999-206147	19990721

Patent Number (1)
JP 2001035664

Kind Code (1)
A

Patent Publication Date (1)
20010209

Application Number (1)
JP 1999-206147

Application Date (1)
19990721

Priority Patent Number (1)
JP 1999-206147

Priority Patent Publication Date (1)
19990721

Abstract The invention relates to an organic electroluminescent device comprising an azaindolizine-containing layer placed between a pair of electrodes.

=> s 14 not 15
L6 234 L4 NOT L5

```

=> s LUMIN? OR ELECTROLUMIN? OR ORGANOLUMIN? OR (ELECTRO OR ORGANO OR ORG#) (2A) LUMIN? OR
LIGHT?(2A) (EMIT? OR EMISSION?) OR EL OR E(W)L OR L(W)E(W)D OR OLED OR DIODE# OR SEMICONDUCTOR#
TRANSISTOR?
343037 LUMIN?
77576 ELECTROLUMIN?
55 ORGANOLUMIN?
89987 ELECTRO
9 ELECTROS
89995 ELECTRO
(ELECTRO OR ELECTROS)
19764 ORGANO
3 ORGANOS
19767 ORGANO
(ORGANO OR ORGANOS)
1060191 ORG#
343037 LUMIN?
13165 (ELECTRO OR ORGANO OR ORG#) (2A) LUMIN?
1256968 LIGHT?
252960 EMIT?
584730 EMISSION?
89593 LIGHT?(2A) (EMIT? OR EMISSION?)
27002 EL
956 ELS
27927 EL
(EL OR ELS)
2090803 E
1625990 L
2064 E(W)L
1625990 L
2090803 E
2536233 D
29 L(W)E(W)D
5184 OLED
2548 OLEDs
6488 OLED
(OLED OR OLEDs)
118570 DIODE#
647637 SEMICONDUCTOR#
151037 TRANSISTOR?
1782 SEMICONDUCTOR# TRANSISTOR?
(SEMICONDUCTOR# (W) TRANSISTOR?)
L7 527912 LUMIN? OR ELECTROLUMIN? OR ORGANOLUMIN? OR (ELECTRO OR ORGANO
OR ORG#) (2A) LUMIN? OR LIGHT?(2A) (EMIT? OR EMISSION?) OR EL OR
E(W)L OR L(W)E(W)D OR OLED OR DIODE# OR SEMICONDUCTOR# TRANSISTO
R?

=> s 16 and 17
L8 7 L6 AND L7

=> d 18 1-7 ibib abs hitstr

```

L8 ANSWER 1 OF 7 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Compositions, methods and kits pertaining to luminescent compounds

Author/Inventor

Wood, Keith; Hawkins, Erika; Scurria, Mike; Klaubert, Dieter

Patent Assignee/Corporate Source

Promega Corporation, USA

Source

PCT Int. Appl., 60 pp. CODEN: PIXXD2

Document Type

Patent

Language

English

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003040100	A1	20030515	WO 2002-US34972	20021101

Patent Number (1)

WO 2003040100

Kind Code (1)

A1

Patent Publication Date (1)

20030515

Application Number (1)
WO 2002-US34972
Application Date (1)
20021101

Priority Patent Number (1)
US 2001-53482

Priority Kind Code (1)
A

Priority Patent Publication Date (1)
20011102

Abstract

A method of measuring the enzymic activity of a luciferase includes contacting a luminogenic protein, such as a luciferase, with a protected luminophore to form a composition; and detecting light produced from the composition. The protected luminophore provides increased stability and improved signal-to-background ratios relative to the corresponding unmodified coelenterazine.

L8 ANSWER 2 OF 7 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Coelenterazine derivatives for improved solution solubility

Author/Inventor

Hawkins, Erika M.; O'Grady, Michael; Klaubert, Dieter; Scurria, Michael; Good, Troy; Stratford, Cathy; Flemming, Rod; Simpson, Dan; Wood, Keith V.

Patent Assignee/Corporate Source

Promega Corporation, Madison, WI, 53715, USA

Source

Bioluminescence & Chemiluminescence: Progress & Current Applications. [Proceedings of the Symposium on Bioluminescence and Chemiluminescence], 12th, Cambridge, United Kingdom, Apr. 5-9, 2002 (2002), 149-152. Editor(s): Stanley, Philip E.; Kricka, Larry J. World Scientific Publishing Co. Pte. Ltd.: Singapore, Singapore. CODEN: 69DPGZ; ISBN: 981-238-156-2

Document Type

Conference

Language

English

Abstract

Intracellular luminescent techniques requiring coelenterazine, such as bioluminescence resonance energy transfer (BRET), calcium detection, and intracellular reporter measurements, must accommodate the poor stability of this substrate in physiolo. buffered solns. Coelenterazine degradation leads both to loss of luminescence over time, and increased background luminescence caused by enzyme-independent oxidation (autoluminescence). Both conditions limit luminescence sensitivity by reducing the signal-to-noise ratio. Coelenterazine can be stabilized by derivatizing the enol oxygen with an acyl oxymethyl ether. This prevents spontaneous oxidation in solution while making the substrate available intracellularly upon cleavage of the blocking group by endogenous esterases. We will describe the stability of pivaloyl oxymethyl coelenterazine-h (POM coelenterazine-h), and the effect of POM coelenterazine-h on intracellular luminescence, autoluminescence, and luminescent reaction kinetics. Also, we will present the characteristics of two other coelenterazine derivs.

L8 ANSWER 3 OF 7 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Preparation of luciferin derivatives of Umihotaru (Cypridina hilgendorffii)

Author/Inventor

Mitani, Motohiro; Sakaki, Hidejiro; Koinuma, Yasuyoshi; Totani, Yoshiaki

Patent Assignee/Corporate Source

Nippon Oils & Fats Co., Ltd., Japan; NOF Corporation

Source

Jpn. Kokai Tokkyo Koho, 8 pp. CODEN: JKXXAF

Document Type

Patent

Language

Japanese

Patent Information

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 08059686	A	19960305	JP 1994-198770	19940823

Patent Number (1)

JP 08059686

Kind Code (1)

A

Patent Publication Date (1)

19960305

Application Number (1)

JP 1994-198770

Application Date (1)

19940823

Priority Patent Number (1)

JP 1994-198770

Priority Patent Publication Date (1)

19940823

Abstract

The title compds. (I; R1, R2 = H, C1-20 alkyl, C6-20 aryl, C7-19 arylalkyl; R3 = C1-5 alkyl or alkoxy; n = 0-5), which are useful as substrates for luminescent determination of sugar hydrolases such as α -D-galactosidase, are prepared by reacting imidazopyrazinone derivs. (II; R1 - R3, n = same as above) with sugar derivs. (III; X = halo; R4 = C1-7 acyl) in the presence of silver triflate and Na2HPO4, followed by solvolysis in the presence of an alkali. Thus, 0.1 g 6-(4-methoxyphenyl)-2-methylimidazo[1,2-*a*]pyrazin-3-one and 1.1 g Na2HPO4 were treated with 5 mL MeCN, 9 mL benzene, and 2.6 g mol. sieve 4A and stirred at room temperature for 1 h, treated with 0.18 g 2,3,4,6-tetra-O-acetyl- α -D-galactopyranosyl bromide and 0.37 g silver triflate, and stirred at room temperature for 2 h to give 39% 6-(4-methoxyphenyl)-2-methyl-3-(2,3,4,6-tetra-O-acetyl- α -D-galactopyranosyloxy)imidazo[1,2-*a*]pyrazine, which (0.5 g) was treated with 3.5 mL MeOH and 1.8 mL concentrated aqueous NH3 and stirred at 40° for 6 h 30 min to give 78% 6-(4-methoxyphenyl)-2-methyl-3-(α -D-galactopyranosyloxy)imidazo[1,2-*a*]pyrazine (IV). IV showed luminescence in the presence of β -D-galactosidase with correlation factor $r = 0.992$.

L8 ANSWER 4 OF 7 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Enhancement effect of 2,6-O-dimethyl- β -cyclodextrin on the chemiluminescent detection of β -D-galactosidase using a Cypridina luciferin analog

Author/Inventor

Mitani, Motohiro; Sakaki, Syujiro; Koinuma, Yasumi; Toya, Yoshiaki; Kosugi, Masanori

Patent Assignee/Corporate Source

Tsukuba Res. Lab., NOF Corp., Tsukuba, 300-26, Japan

Source

Analytical Sciences (1995), 11(6), 1013-15 CODEN: ANSCEN; ISSN: 0910-6340

Document Type

Journal

Language

English

Abstract

β -Cyclodextrins enhanced the chemiluminescent detection of β -galactosidase using the Cypridina luciferin analog 3-(β -D-galactopyranosyloxy)-6-(4-methoxyphenyl)-2-methylimidazo[1,2-*a*]pyrazine (β -Gal-MCLA) in the order 2,6-O-dimethyl- β -cyclodextrin > 2,3,6-O-trimethyl- β -cyclodextrin > β -cyclodextrin. Detection of mouse IgG by chemiluminescent enzyme immunoassay (CLEIA) using β -Gal-MCLA and β -galactosidase to amplify the signal was also enhanced by inclusion of 2,6-O-trimethyl- β -cyclodextrin.

L8 ANSWER 5 OF 7 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Mechanism of photoinactivation and re-activation in the bioluminescence system of the ctenophore Mnemiopsis

Author/Inventor

Antcil, Michel; Shimomura, Osamu

Patent Assignee/Corporate Source

Mar. Biol. Lab., Woods Hole, MA, 02543, USA

Source

Biochemical Journal (1984), 221(1), 269-72 CODEN: BJQOAK; ISSN: 0306-3275

Document Type

Journal

Language

English

Abstract

The bioluminescence of *M. leidyi* takes place when the photoprotein mnemiopsin in the photocytes reacts with Ca^{2+} . The luminescence is inhibited in sunlight and this photoinhibition is reversible by keeping the live specimens in the dark. Exts. of mnemiopsin are similarly photoinhibited, but the photoinhibition cannot be reversed in the dark. Photoinhibited mnemiopsin can be reactivated in the dark by incubation with coelenterazine and O only in solns. having a pH very close to 9.0. The reactivation in vivo probably takes place in the same manner, using the coelenterazine that is supplied from its abundant storage form. Apparently, photoinactivation of mnemiopsin results in the dissociation of coelenterazine and O from the mol. of photoprotein; the dissociated form of the former mol. is an inactive form of coelenterazine, not free coelenterazine.

L8 ANSWER 6 OF 7 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Comparison of the amounts of key components in the bioluminescence systems of various coelenterates

Author/Inventor

Shimomura, Osamu; Johnson, Frank H.

Patent Assignee/Corporate Source

Dep. Biol., Princeton Univ., Princeton, NJ, 08540, USA

Source

Comparative Biochemistry and Physiology, Part B: Biochemistry & Molecular Biology (1979), 64B(1), 105-7 CODEN: CBPBB8; ISSN: 0305-0491

Document Type

Journal

Language

English

Abstract

Luciferase, photoprotein, free and protein-bound coelenterazine (l) and l enol-sulfate were assayed and compared in 5 bioluminescent coelenterates. Hydrozoans Aequorea aequorea and Halistaura cellularia contained photoprotein plus very small amts. of l enol-sulfate and luciferase activity, but no free l. Anthozoans Ptilosarcus gurneyi, Cavemularia obesa, and Renilla muelleri contained luciferase, l, and l enol-sulfate, but very little or no photoprotein. l existed mainly in a stabilized form bound to a Ca-binding protein. The bioluminescent reactions in the coelenterates were compared.

L8 ANSWER 7 OF 7 CAPLUS COPYRIGHT 2008 ACS on STN

Title

Chemical nature of bioluminescence systems in coelenterates

Author/Inventor

Shimomura, Osamu; Johnson, Frank H.

Patent Assignee/Corporate Source

Dep. Biol., Princeton Univ., Princeton, NJ, USA

Source

Proceedings of the National Academy of Sciences of the United States of America (1975), 72(4), 1546-9 CODEN: PNASA6; ISSN: 0027-8424

Document Type

Journal

Language

English

Abstract

Anal. of substances involved in light -emitting reactions among bioluminescent coelenterates revealed a pronounced uniformity in the structural features of initial reactants, i.e., luciferins and photoprotein chromophores, as well as the light - emitter product. This product is structurally identical among the different classes of coelenterates: i.e., Hydrozoa (the jellyfish, Aequorea), Anthozoa (the sea cactus, Cavemularia; sea pansy, Renilla; and sea pen, Leioptilus), and very likely also the Scyphozoa (the jellyfish, Pelagia). In each of these instances the reaction product, 2-(p-hydroxyphenylacetyl)amino-3-benzyl-5-(p-hydroxyphenyl) pyrazine, is the actual light - emitter, whether it occurs in a Ca2+-triggered photoprotein type of luminescence or in a luciferin-luciferase type. The evidence indicates that in certain coelenterates, e.g., Cavemularia, these 2 types are equally significant, whereas in others (Renilla and Leioptilus) the luciferin-luciferase type predominates over the Ca-triggerable photoprotein type. Only the photoprotein type functions in the luciferaseless jellyfish, Aequorea. In all instances investigated, the structure of the light - emitter prior to the luminescence reaction appears to be essentially the same as that of the chromophore of unreacted aequorin. The product of the luminescence reaction is absent in exts. of nonluminous species. However, a product very similar to that of luminescent coelenterates occurs also in representatives of other phyla, including the cephalopod molluscs, e.g., the "firefly squid" Watasenia and probably various ctenophores as well.

-->

-->

-->